

09/787083

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**BOX PCT/DO-EO**

Commissioner for Patents
Washington, D.C. 20231

"Express Mail" mailing label number **EF 391 026 502 US**

I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 CFR § 1.10 on the date indicated above and addressed to: Commissioner of Patents, Washington, DC 20231, on March 14, 2001.


Colleen Hanagan

NATIONAL STAGE APPLICATION TRANSMITTAL LETTER
APPLICATION FILING UNDER 35 U.S.C. § 371

Transmitted herewith for filing is the patent application of.

Inventor(s)/Applicant(s):	Ruelle, Jean-Louis
International Application No.:	PCT/EP99/06781
International Published Appln. No.:	WO 00/15802
International Filing Date:	14 September 1999
Priority Filing Date:	14 August 1998
Thirty Month Date:	14 March 2001
Title:	"MORAXELLA CATARRHALIS BASB034 POLYPEPTIDES AND USES THEREOF"

1. **THIS NEW APPLICATION IS A NATIONAL STAGE APPLICATION UNDER PCT, CHAPTER II WITH A REQUEST FOR EXAMINATION WITHOUT DELAY TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US).**

- ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. § 371,
☐ This is a **SECOND** or subsequent submission of items concerning a filing under 35 U.S.C. § 371.

2. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).

104150 2802650

09/787083

PATENT

DOCKET NO. BM45332

JC08 Rec'd PCT/PTO 14 MAR 2001

3. A proper Demand for International Preliminary Examination was made by the 19TH month from the earliest claimed priority date.

4. Enclosed items are required for filing under 37 CFR § 1.53(b) and § 1.494(b) or § 1.495(b):

- ☒ One copy of International Publication **WO 00/15802**
- (a) ☒ is transmitted herewith (**required only if not transmitted by the International Bureau**)
- (b) ☒ has been transmitted by the International Bureau
- (c) ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)

Fees

☒ The basic national fee set forth in 37 CFR § 1.482 - International Preliminary Examination Fee not paid to USPTO but International Search Report prepared by the EPO or JPO - **\$860.00**

- ☒ Claims in Excess of 20 (13 @ \$18.00)
- ☒ Independent Claims in Excess of 3 (2 @ \$80.00)

5. Further enclosed are:

- ☒ One copy of International Preliminary Examination Report.
- ☒ One copy of International Search Report - (references not available).
- ☐ One copy of Written Opinion.
- ☐ One copy of PCT Request as filed.
- ☐ One copy of Chapter II Demand as filed.

6. ☐ A translation of the International Application into English (35 U.S.C. § 371(c)(2))

7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. § 371(c)(3))

8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. § 371(c)(3))

9. Still additional papers enclosed:

- ☐ Assignment with Assignment Recordation Form Cover Sheet
- ☐ Verification Statement Claiming Small Entity Status
- ☐ Declaration or oath is enclosed executed by the inventor
- ☐ An Information Disclosure Statement under 37 CFR § 1.97 and § 1.98
- ☒ Return Acknowledgment Postcard

10. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)

A. Enclosed are:

- (a) ☒ Computer Readable Copy of the Sequence Listing
- (b) ☒ Paper Copy (identical to Computer Readable Copy) of the Sequence Listing

B. ☐ Enclosed is a paper copy of the Sequence Listing. This paper copy and a Computer Readable Form thereof are identical with the Computer Readable Form in another application of the Applicant which is fully identified as follows.

U.S. Application No.: @@

Filed: @@

Attorney Docket No.: @@

which is believed to comply with the rules set forth in 37 CFR § 1.821 et. seq. Applicants requests pursuant to 37 CFR § 1.821(e) that this Computer Readable Form be used in the present application. **Please TRANSFER the sequence listing from the parent to this application.**

C. ☒ Statement under 37 CFR § 1.821(f). **The information recorded in computer readable form is identical to the written Sequence Listing.**D. ☒ Statement under 37 CFR § 1.821(g) (required when Sequence Listing not submitted at the time of filing under 35 U.S.C. § 111(a)) or 37 CFR § 1.821(f) (required when Sequence Listing not submitted at the time of filing under the Patent Cooperation Treaty): **The submission of the Sequence Listing includes no new matter.**E. ☒ Amendment. Please enter the Sequence Listing into the application11. **Preliminary Amendment**

Prior to calculation of fees, kindly enter

- ☒ Preliminary Amendment submitted herewith
- ☐ do not enter Preliminary Amendment

09/787083

PATENT

DOCKET NO. BM45332

JCOG Rec'd PGT/PTO 14 MAR 2001


12. Fee payment being made at this time is enclosed:

* Basic filing fee (\$860.00)	860 00
* Claims in Excess of 20	234.00
* (13 @ \$18 00)	
Independent Claims in Excess of 2	160.00
(2 @ \$80.00)	
* Total Fees enclosed:	<u>\$1254 00</u>

13. The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Account No 50-0258. This letter is filed in duplicate for accounting purposes.

Respectfully submitted,

Date: March 13, 2001



Arthur E. Jackson

Registration No. 34,354

Allen Bloom

Registration No. 29,135

Attorney for Applicant

DECHERT

4000 Bell Atlantic Tower

1717 Arch Street

Philadelphia, PA 19103-2793

Fax: (609) 620-3259

Attn: Arthur E. Jackson, Esq.
(609) 620-3254

107273 1 07

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Ruelle
Serial No.: Unknown
Filed: Herewith

Group Art Unit No.: Unknown
Examiner: Unknown

For: *Moraxella Catarrhalis* BASB034 Polypeptides and Uses Thereof

PRELIMINARY AMENDMENT

Sir:

Please enter this Preliminary Amendment into the record of the above-identified patent application before the calculation of fees.

In the Claims:

Please delete the claims of the application as filed in the PCT and substitute therefor:

27. An isolated polypeptide comprising a member selected from the group consisting of
- (a) an amino acid sequence which has at least 90% identity to SEQ ID NOs:2, 4, 6, or 8;
 - (b) an immunogenic fragment of the amino acid sequence of (a), wherein the immunogenic fragment is at least 90% identical to an aligned contiguous segment of SEQ ID NOs:2, 4, 6, or 8; and
 - (c) an immunogenic fragment of the amino acid sequence of (a) that matches an aligned contiguous segment of SEQ ID NOs:2, 4, 6, or 8 with no more than five single amino acid substitutions, deletions or additions;

wherein the isolated polypeptide, when administered to a subject in a suitable composition which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an immune response that recognizes a polypeptide having the sequence of SEQ ID NOs:2, 4, 6, or 8.

28. The isolated polypeptide of claim 27, wherein the immunogenic fragment of (b) comprises at least 15 amino acids.

29. The isolated polypeptide of claim 28, wherein the immunogenic fragment matches an aligned contiguous segment of SEQ ID NOs:2, 4, 6, or 8 with no more than a single amino acid substitution, deletion or addition.

30. The isolated polypeptide of claim 27, wherein the immunogenic fragment of (b) comprises at least 20 amino acids.
31. The isolated polypeptide of Claim 27 wherein the amino acid sequence of (a) has at least 95% identity to SEQ ID NOs:2, 4, 6, or 8.
32. The isolated polypeptide of Claim 31 wherein the isolated polypeptide comprises the amino acid sequence of SEQ ID NOs:2, 4, 6, or 8.
33. The isolated polypeptide of claim 31 wherein the isolated polypeptide consists of the amino acid sequence of SEQ ID NOs:2, 4, 6, or 8.
34. A fusion protein comprising the isolated polypeptide of Claim 27.
35. The isolated polypeptide of Claim 27 wherein the polypeptide is the immunogenic fragment having no more than two single amino acid substitutions, deletions or additions relative to the aligned sequence.
36. The isolated polypeptide of Claim 27 wherein the polypeptide is the immunogenic fragment having no more than one single amino acid substitution, deletion or addition relative to the aligned sequence.
37. The isolated polypeptide of Claim 27 wherein the polypeptide is the immunogenic fragment which matches the aligned sequence.
38. An isolated polypeptide encoded by an isolated first polynucleotide wherein the isolated first polynucleotide hybridizes under stringent conditions to a second polynucleotide which encodes the polypeptide of SEQ ID NOs:2, 4, 6, or 8; wherein stringent conditions comprise overnight incubation at 42° C in a solution comprising: 50% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5× Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1× SSC at about 65° C; wherein the isolated polypeptide,

when administered to a subject in a suitable composition which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an immune response that recognizes a polypeptide having the sequence of SEQ ID NOs:2, 4, 6, or 8.

39. An isolated polynucleotide encoding a polypeptide of Claim 27 or the full complement to the isolated polynucleotide.

40. An isolated polynucleotide encoding a polypeptide of Claim 27, wherein the isolated polynucleotide encodes the polypeptide comprising SEQ ID NOs:2, 4, 6, or 8.

41. An isolated polynucleotide comprising the polynucleotide of SEQ ID NOs:1, 3, 5, or 7.

42. An isolated polynucleotide segment comprising a polynucleotide sequence or the full complement of the entire length of the polynucleotide sequence, wherein the polynucleotide sequence hybridizes to the full complement of SEQ ID NOs:1, 3, 5, or 7 minus the complement of any stop codon, wherein the hybridization conditions include incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing in 0.1x SSC at 65°C; and, wherein the polynucleotide sequence is identical to SEQ ID NOs:1, 3, 5, or 7 minus any terminal stop codon, except that, over the entire length corresponding to SEQ ID NOs:1, 3, 5, or 7 minus any terminal stop codon, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression

$$n_n \leq x_n - (x_n \bullet y)$$

wherein x_n is the total number of nucleotides in SEQ ID NOs:1, 3, 5, or 7 minus any terminal stop codon, y is at least 0.95, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n ; and wherein the polynucleotide sequence detects *Moraxella catarrhalis*.

43. An expression vector comprising the isolated polynucleotide of Claim 39.

44. A host cell transformed with the expression vector of Claim 43.

45. A process of producing an isolated polypeptide comprising (a) culturing the host cell of Claim 44 under conditions sufficient for the production of the encoded polypeptide and (b) recovering the polypeptide.

46. A nucleic acid vaccine comprising the isolated polynucleotide of Claim 39 and a pharmaceutically acceptable carrier.

47. An isolated polynucleotide segment comprising a polynucleotide sequence or the full complement of the entire length of the polynucleotide sequence, wherein the polynucleotide sequence is identical to SEQ ID NOs:1, 3, 5, or 7 minus any terminal stop codon, except that, over the entire length corresponding to SEQ ID NOs:1, 3, 5, or 7 minus any terminal stop codon, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression

$$n_n \leq x_n - (x_n \bullet y)$$

wherein x_n is the total number of nucleotides in SEQ ID NOs:1, 3, 5, or 7 minus any terminal stop codon, y is at least 0.90, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n ; and wherein the polynucleotide sequence detects *Moraxella catarrhalis*.

48. The isolated polynucleotide of Claim 47 where y is at least 0.95.

49. An expression vector comprising the isolated polynucleotide of Claim 47 which codes for a polypeptide that, when administered to a mammal which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an immune response that recognizes a polypeptide having the sequence of SEQ ID NOs:2, 4, 6, or 8.

50. A host cell transformed with the isolated polynucleotide or an expression vector comprising the isolated polynucleotide of Claim 47.

51. A process of producing an isolated polypeptide comprising (a) culturing the host cell of Claim 50 under conditions sufficient for the production of the encoded polypeptide and (b) recovering the polypeptide.

52. A vaccine comprising the polypeptide of Claim 27 and a pharmaceutically acceptable carrier.
53. The vaccine of Claim 52, wherein the composition comprises at least one other *Moraxella catarrhalis* antigen.
54. An antibody immunospecific for the polypeptide or immunogenic fragment of Claim 27.
55. A method for inducing an immune response in a mammal comprising administration of the polypeptide of Claim 27.
56. A method of diagnosing a *Moraxella catarrhalis* infection, comprising identifying a polypeptide of Claim 27, or an antibody that is immunospecific for the polypeptide, present within a biological sample from an animal suspected of having such an infection.
57. A method for inducing an immune response in a mammal comprising administration of the isolated polynucleotide of Claim 39.
58. A therapeutic composition useful in treating humans with *Moraxella catarrhalis* comprising at least one antibody directed against the polypeptide of claim 27 and a suitable pharmaceutical carrier.
59. A process for expressing the polynucleotide of Claim 39 comprising transforming a host cell with the expression vector comprising the polynucleotide and culturing the host cell under conditions sufficient for expression of the polynucleotide.

REMARKS

Applicant respectfully requests that this Preliminary Amendment be entered in this case before the calculation of fees and before examination of the subject application.

Claims

Claims 1-26 have been canceled without prejudice or disclaimer of the subject matter therein. Applicant reserves the right to prosecute, in one or more patent applications, the canceled claims, the claims to non-elected inventions, the claims as originally filed, and any other claims supported by the specification.

New claims 27-59 have been introduced. No new matter is added.

Support


Support for the new claims is either obvious, or is as described in the text below. Particularly, support for the recitation of “five single amino acid substitutions, deletions or additions” may be found, for example, at page 7, lines 4-5. Support for compositions of the isolated polypeptide which include an adjuvant recited in the claims may be found, for example, at page 41, lines 1-2. Support for the stringent hybridization conditions may be found, for example, at page 14, line 28 through page 15, line 4. Support for the recitation of sequence relatedness such as in claim 47 may be found in the specification, for example, at page 44, line 12 through page 45, line 8.

International Application No.: PCT/EP99/06781
Attorney Docket No.: BM45332

Closing Remarks

Entry of this Preliminary Amendment prior to calculation of fees and Examination and Allowance of the pending claims are respectfully requested.

Respectfully submitted,



Arthur E. Jackson
Registration No. 34,354
Allen Bloom
Registration No. 29,135
Attorneys for Applicant

DECHERT PRICE & RHOADS
Princeton Pike Corporate Center
PO Box 5218
Princeton, New Jersey 08543-5218
Allen Bloom (609) 620-3214
Arthur E. Jackson (609) 620-3254
Fax: (609) 620-3259
Attn: Allen Bloom, Esq.
(609) 620-3214

MORAXELLA CATARRHALIS BASB034 POLYPEPTIDES AND USES THEREOF**FIELD OF THE INVENTION**

This invention relates to polynucleotides, (herein referred to as "BASB034 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB034" or "BASB034 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the invention relates to diagnostic assays for detecting infection of certain pathogens.

BACKGROUND OF THE INVENTION

Moraxella catarrhalis (also named *Branhamella catarrhalis*) is a Gram negative bacteria frequently isolated from the human upper respiratory tract. It is responsible for several pathologies the main ones being otitis media in infants and children, and pneumonia in elderlies. It is also responsible of sinusitis, nosocomial infections and less frequently of invasive diseases.

Otitis media is an important childhood disease both by the number of cases and its potential sequelae. More than 3.5 millions cases are recorded every year in the United States, and it is estimated that 80 % of the children have experienced at least one episode of otitis before reaching the age of 3 (Klein, JO (1994) Clin.Inf.Dis 19:823). Left untreated, or becoming chronic, this disease may lead to hearing losses that could be temporary (in the case of fluid accumulation in the middle ear) or permanent (if the auditive nerve is damaged). In infants, such hearing losses may be responsible for a delayed speech learning.

Three bacterial species are primarily isolated from the middle ear of children with otitis media: *Streptococcus pneumoniae*, non typeable *Haemophilus influenza* (NTHi) and *M. catarrhalis*. They are present in 60 to 90 % of the cases. A review of recent studies shows that *S. pneumoniae* and NTHi represent both about 30 %. and *M. catarrhalis* about 15 % of

the otitis media cases (Murphy, TF (1996) Microbiol.Rev. 60:267). Other bacteria could be isolated from the middle ear (*H. influenza* type B, *S. pyogenes* etc) but at a much lower frequency (2 % of the cases or less).

- 5 Epidemiological data indicate that, for the pathogens found in the middle ear, the colonization of the upper respiratory tract is an absolute prerequisite for the development of an otitis; other are however also required to lead to the disease (Dickinson, DP et al. (1988) J. Infect.Dis. 158:205, Faden, HL et al. (1991) Ann.Otorhinol.Laryngol. 100:612). These are important to trigger the migration of the bacteria into the middle ear via the Eustachian
10 tubes, followed by the initiation of an inflammatory process. These factors are unknown todate. It has been postulated that a transient anomaly of the immune system following a viral infection, for example, could cause an inability to control the colonization of the respiratory tract (Faden, HL et al (1994) J. Infect.Dis. 169:1312). An alternative explanation is that the exposure to environmental factors allow a more important colonization of some
15 children, who subsequently become susceptible to the development of otitis media because of the sustained presence of middle ear pathogens (Murphy, TF (1996) Microbiol.Rev. 60:267).

- The immune response to *M. catarrhalis* is poorly characterized. The analysis of strains
20 isolated sequentially from the nasopharynx of babies followed from 0 to 2 years of age, indicates that they get and eliminate frequently new strains. This indicates that an efficacious immune response against this bacteria is mounted by the colonized children (Faden, HL et al (1994) J. Infect.Dis. 169:1312).

- 25 In most adults tested, bactericidal antibodies have been identified (Chapman, AJ et al. (1985) J. Infect.Dis. 151:878). Strains of *M. catarrhalis* present variations in their capacity to resist serum bactericidal activity: in general, isolates from diseased individuals are more resistant than those who are simply colonized (Hol. C et al. (1993) Lancet 341:1281. Jordan. KL et al. (1990) Am.J.Med. 88 (suppl. 5A):28S). Serum resistance could therefore be

considered as a virulence factor of the bacteria. An opsonizing activity has been observed in the sera of children recovering from otitis media.

The antigens targetted by these different immune responses in humans have not been

5 identified, with the exception of OMP B1, a 84 kDa protein which expression is regulated by iron, and that is recognized by the sera of patients with pneumonia (Sethi, S, et al. (1995) Infect.Immun. 63:1516) , and of UspA1 and UspA2 (Chen D. et al.(1999), Infect.Immun. 67:1310).

10 A few other membrane proteins present on the surface of *M. catarrhalis* have been characterized using biochemical method, or for their potential implication in the induction of a protective immunity (for review, see Murphy, TF (1996) Microbiol.Rev. 60:267). In a mouse pneumonia model, the presence of antibodies raised against some of them (UspA, CopB) favors a faster clearance of the pulmonary infection. Another polypeptide (OMP CD)
15 is highly conserved among *M. catarrhalis* strains, and presents homologies with a porin of *Pseudomonas aeruginosa*, which has been demonstrated efficacious against this bacterium in animal models.

The frequency of *Moraxella catarrhalis* infections has risen dramatically in the past few
20 decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Moraxella catarrhalis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this
25 organism.

SUMMARY OF THE INVENTION

The present invention relates to BASB034, in particular BASB034 polypeptides and BASB034 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including prevention and treatment of microbial diseases, amongst others.

- 5 In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting expression or activity of BASB034 polynucleotides or polypeptides.
- 10 Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

- 15 The invention relates to BASB034 polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of BASB034 of *Moraxella catarrhalis*, which is related by amino acid sequence homology to *Klebsiella pneumoniae* outer membrane phospholipase A protein. The invention relates especially to BASB034 having the nucleotide and amino acid sequences set out in SEQ ID
- 20 NO:1, 3, 5 or 7 and SEQ ID NO:2, 4, 6 or 8 respectively. It is understood that sequences recited in the Sequence Listing below as "DNA" represent an exemplification of one embodiment of the invention, since those of ordinary skill will recognize that such sequences can be usefully employed in polynucleotides in general, including ribopolynucleotides.

25

Polypeptides

In one aspect of the invention there are provided polypeptides of *Moraxella catarrhalis* referred to herein as "BASB034" and "BASB034 polypeptides" as well as biologically,

diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

The present invention further provides for:

- 5 (a) an isolated polypeptide which comprises an amino acid sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% or exact identity, to that of SEQ ID NO:2, 4, 6 or 8;
- (b) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more
- 10 preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1, 3, 5 or 7 over the entire length of SEQ ID NO:1, 3, 5 or 7 respectively; or
- (c) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or
- 15 exact identity, to the amino acid sequence of SEQ ID NO:2, 4, 6 or 8.

The BASB034 polypeptides provided in SEQ ID NO:2, 4, 6 or 8 are the BASB034 polypeptides from *Moraxella catarrhalis* strains Mc2931 (ATCC 43617), Mc2908, Mc2913 and Mc2969.

20

- The invention also provides an immunogenic fragment of a BASB034 polypeptide, that is, a contiguous portion of the BASB034 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2, 4, 6 or 8; That is to say, the fragment (if necessary when coupled to a
- 25 carrier) is capable of raising an immune response which recognises the BASB034 polypeptide. Such an immunogenic fragment may include, for example, the BASB034 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB034 according to the invention comprises substantially all of the extracellular

domain of a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2, 4, 6 or 8 over the entire length of SEQ ID NO:2

- 5 A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with BASB034 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

10

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2, 4, 6 or 8 or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence.

- 15 Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

20

Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2, 4, 6 or 8, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO:2, 4, 6 or 8.

25

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments

may be employed as intermediates for producing the full-length polypeptides of the invention.

Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids
5 are substituted, deleted, or added in any combination.

The polypeptides, or immunogenic fragments, of the invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which
10 contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

15 In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of
20 the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

Furthermore, this invention relates to processes for the preparation of these fusion
25 proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared to non-fused protein. The fusion partner may assist in providing T helper epitopes (immunological fusion partner), preferably T helper epitopes recognised by humans, or
5 assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

Fusion partners include protein D from *Haemophilus influenzae* and the non-structural
10 protein from influenzae virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesize an N-acetyl-L-alanine amidase, amidase LytA, (coded by the lytA gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The
15 C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of E.coli C-LytA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible
20 to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188 - 305.

The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions,
25 whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr.

Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well
5 understood in the art.

It is most preferred that a polypeptide of the invention is derived from *Moraxella catarrhalis*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from
10 organisms of the same taxonomic family or order.

Polynucleotides

It is an object of the invention to provide polynucleotides that encode BASB034 polypeptides, particularly polynucleotides that encode the polypeptide herein designated
15 BASB034.

In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB034 polypeptides comprising a sequence set out in SEQ ID NO:1, 3, 5 or 7 which includes a full length gene, or a variant thereof.
20

The BASB034 polynucleotides provided in SEQ ID NO:1, 3, 5 or 7 are the BASB034 polynucleotides from *Moraxella catarrhalis* strains Mc2931 (ATCC 43617), Mc2908, Mc2913 and Mc2969.

25 As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB034 polypeptides and polynucleotides, particularly *Moraxella catarrhalis* BASB034 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically,

diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

Another aspect of the invention relates to isolated polynucleotides, including at least one full
5 length gene, that encodes a BASB034 polypeptide having a deduced amino acid sequence of SEQ ID NO:2, 4, 6 or 8 and polynucleotides closely related thereto and variants thereof.

In another particularly preferred embodiment of the invention there is a BASB034 polypeptide from *Moraxella catarrhalis* comprising or consisting of an amino acid
10 sequence of SEQ ID NO:2, 4, 6 or 8 or a variant thereof.

Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID NO:1, 3, 5 or 7, a polynucleotide of the invention encoding BASB034 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and
15 sequencing chromosomal DNA fragments from bacteria using *Moraxella catarrhalis* Catlin cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1, 3, 5 or 7, typically a library of clones of chromosomal DNA of *Moraxella catarrhalis* Catlin in *E.coli* or some other suitable host is probed with a radiolabeled
20 oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both
25 directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.: Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By

Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene sequence. Illustrative of the invention, each polynucleotide set out in SEQ ID NO:1, 3, 5 or 7 was discovered in a DNA library derived from *Moraxella catarrhalis*.

5

Moreover, each DNA sequence set out in SEQ ID NO:1, 3, 5 or 7 contains an open reading frame encoding a protein having about the number of amino acid residues set forth in SEQ ID NO:2, 4, 6 or 8 with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art.

10

The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1327 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

The polynucleotide of SEQ ID NO:3, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1327 of SEQ ID NO:3, encodes the polypeptide of SEQ ID NO:4.

The polynucleotide of SEQ ID NO:5, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1327 of SEQ ID NO:5, encodes the polypeptide of SEQ ID NO:6.

The polynucleotide of SEQ ID NO:7, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1327 of SEQ ID NO:7, encodes the polypeptide of SEQ ID NO:8.

In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

- (a) a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1, 3, 5 or 7 over the entire length of SEQ ID NO:1, 3, 5 or 7 respectively; or

(b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2, 4, 6 or 8, over the entire length of SEQ ID NO:2, 4, 6 or 8 respectively.

5

A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Moraxella catarrhalis*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions (for example, using a temperature in the range of 45 – 65°C and an SDS
10 concentration from 0.1 – 1%) with a labeled or detectable probe consisting of or comprising the sequence of SEQ ID NO:1, 3, 5 or 7 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

The invention provides a polynucleotide sequence identical over its entire length to a coding
15 sequence (open reading frame) in SEQ ID NO:1, 3, 5 or 7. Also provided by the invention is a coding sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least
20 one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding
25 additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson *et al.*, *Cell* 37: 767 (1984), both of which may be useful in purifying

polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

- 5 The nucleotide sequence encoding BASB034 polypeptide of SEQ ID NO:2, 4, 6 or 8 may be identical to the polypeptide encoding sequence contained in nucleotides 1 to 1326 of SEQ ID NO:1, 3, 5 or 7 respectively. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2, 4, 6 or 8.

10

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Moraxella catarrhalis* BASB034 having an amino acid sequence set out in SEQ ID NO:2, 4, 6 or 8.

- 15 The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

20

The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2, 4, 6 or 8. Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

25

Further particularly preferred embodiments are polynucleotides encoding BASB034 variants, that have the amino acid sequence of BASB034 polypeptide of SEQ ID NO:2, 4, 6 or 8 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred

among these are silent substitutions, additions and deletions, that do not alter the properties and activities of BASB034 polypeptide.

- Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB034 polypeptide having an amino acid sequence set out in SEQ ID NO:2, 4, 6 or 8, and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 90% identical over its entire length to a polynucleotide encoding BASB034 polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 95% identical over their entire length to the same are particularly preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.
- Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of SEQ ID NO:1, 3, 5 or 7.

- In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB034 polynucleotide sequences, such as those polynucleotides in SEQ ID NO:1, 3, 5 or 7.

- The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x

SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1, 3, 5 or 7 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1, 3, 5 or 7 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB034 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB034 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

A coding region of a BASB034 gene may be isolated by screening using a DNA sequence provided in SEQ ID NO:1, 3, 5 or 7 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then

used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

There are many methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short DNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, *et al.*, *PNAS USA* 85:908-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases, particularly human diseases, as further discussed herein relating to polynucleotide assays.

The polynucleotides of the invention that are oligonucleotides derived from a sequence of SEQ ID NOS:1 - 8 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also

have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff *et al.*, *Hum Mol Genet* (1992) 1: 363, Manthorpe *et al.*, *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu *et al.*, *J Biol Chem.* (1989) 264: 16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda *et al.*, *Science* (1989) 243: 375), particle bombardment (Tang *et al.*, *Nature* (1992) 356:152, Eisenbraun *et al.*, *DNA Cell Biol* (1993) 12: 791) and *in vivo* infection using cloned retroviral vectors (Seeger *et al.*, *PNAS USA* (1984) 81: 5849).

Vectors, Host Cells, Expression Systems

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis, *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, *Neisseria meningitidis* and *Moraxella catarrhalis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from

transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from
5 plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any
10 of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular
15 environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the present invention can be recovered and purified from recombinant
20 cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques
25 for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant

- virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), Listeria, Salmonella, Shigella, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

10 **Diagnostic, Prognostic, Serotyping and Mutation Assays**

- This invention is also related to the use of BASB034 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB034 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB034 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.
- 20 Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the
- 25 same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a

different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled BASB034 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting
5 differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers *et al.*, *Science*, 230: 1242 (1985).
10 Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for example, Cotton *et al.*, *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

In another embodiment, an array of oligonucleotide probes comprising BASB034
15 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee *et al.*, *Science*, 274: 610 (1996)).
20

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:
(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1, 3, 5 or 7, or a fragment thereof;
(b) a nucleotide sequence complementary to that of (a);
25 (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2, 4, 6 or 8 or a fragment thereof; or
(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2, 4, 6 or 8.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a Disease, among others.

- 5 This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferable, SEQ ID NO:1, 3, 5 or 7, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease,
- 10 which results from under-expression, over-expression or altered expression of the polynucleotide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.
- 15 Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such
- 20 as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding BASB034 polypeptide can be used to identify and analyze mutations.

- The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5'
- 25 and/or the 3' end. These primers may be used for, among other things, amplifying BASB034 DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide

sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial
5 infections, more preferably infections caused by *Moraxella catarrhalis*, comprising
determining from a sample derived from an individual, such as a bodily material, an
increased level of expression of polynucleotide having a sequence of SEQ ID NO:1, 3, 5
or 7. Increased or decreased expression of a BASB034 polynucleotide can be measured
using any one of the methods well known in the art for the quantitation of polynucleotides,
10 such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting,
spectrometry and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-
expression of BASB034 polypeptide compared to normal control tissue samples may be
15 used to detect the presence of an infection, for example. Assay techniques that can be used
to determine levels of a BASB034 polypeptide, in a sample derived from a host, such as a
bodily material, are well-known to those of skill in the art. Such assay methods include
radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich
assays, antibody detection and ELISA assays.

20

The polynucleotides of the invention may be used as components of polynucleotide
arrays, preferably high density arrays or grids. These high density arrays are
particularly useful for diagnostic and prognostic purposes. For example, a set of spots
each comprising a different gene, and further comprising a polynucleotide or
25 polynucleotides of the invention, may be used for probing, such as using hybridization
or nucleic acid amplification, using a probes obtained or derived from a bodily sample,
to determine the presence of a particular polynucleotide sequence or related sequence in
an individual. Such a presence may indicate the presence of a pathogen, particularly
Moraxella catarrhalis, and may be useful in diagnosing and/or prognosing disease or a

course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1, 3, 5 or 7 are preferred. Also preferred is a comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2, 4, 6 or 8.

5

Antibodies

The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

10

In certain preferred embodiments of the invention there are provided antibodies against BASB034 polypeptides or polynucleotides.

15

Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

20

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

25

Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-BASB034 or from naive libraries (McCafferty, *et al.*, (1990), *Nature* 348, 552-554; Marks, *et al.*,
5 (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson *et al.*, (1991) *Nature* 352: 628).

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or
10 polynucleotides by, for example, affinity chromatography.

Thus, among others, antibodies against BASB034-polypeptide or BASB034-polynucleotide may be employed to treat infections, particularly bacterial infections.

15 Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be
20 "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

25

Antagonists and Agonists - Assays and Molecules

Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural

substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan *et al.*, *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

5 The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB034 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB034 polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and BASB034 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and and/or functionally related polypeptides (see D. Bennett *et al.*, *J Mol Recognition*, 8:52-58 (1995); and K. Johanson *et al.*, *J Biol Chem.* 270(16):9459-9471 (1995)).

- The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and/or polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.
- 10 The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of BASB034 polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a
- 15 membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB034 polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a BASB034 agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the BASB034 polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from
- 20 such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of BASB034 polypeptide are most likely to be good antagonists. Molecules that bind well and, as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or
- 25 chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate converted into product, a reporter gene that is responsive to changes in BASB034 polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for BASB034 agonists is a competitive assay that combines BASB034 and a potential agonist with BASB034-binding molecules, recombinant BASB034 binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. BASB034 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB034 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB034-induced activities, thereby preventing the action or expression of BASB034 polypeptides and/or polynucleotides by excluding BASB034 polypeptides and/or polynucleotides from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB034.

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of

various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

- 5 Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

10

Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno

- 15 or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

- The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a
- 20 pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion
- 25 between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB034 proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

In accordance with yet another aspect of the invention, there are provided BASB034 agonists and antagonists, preferably bacteristatic or bactericidal agonists and antagonists.

5 The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

In a further aspect, the present invention relates to mimotopes of the polypeptide of the invention. A mimotope is a peptide sequence, sufficiently similar to the native peptide (sequentially or structurally), which is capable of being recognised by antibodies which
10 recognise the native peptide; or is capable of raising antibodies which recognise the native peptide when coupled to a suitable carrier.

Peptide mimotopes may be designed for a particular purpose by addition, deletion or substitution of elected amino acids. Thus, the peptides may be modified for the purposes
15 of ease of conjugation to a protein carrier. For example, it may be desirable for some chemical conjugation methods to include a terminal cysteine. In addition it may be desirable for peptides conjugated to a protein carrier to include a hydrophobic terminus distal from the conjugated terminus of the peptide, such that the free unconjugated end of the peptide remains associated with the surface of the carrier protein. Thereby
20 presenting the peptide in a conformation which most closely resembles that of the peptide as found in the context of the whole native molecule. For example, the peptides may be altered to have an N-terminal cysteine and a C-terminal hydrophobic amidated tail. Alternatively, the addition or substitution of a D-stereoisomer form of one or more of the amino acids may be performed to create a beneficial derivative, for example to
25 enhance stability of the peptide.

Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number

of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

5 Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB034 polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response

10 to protect said individual from infection, particularly bacterial infection and most particularly *Moraxella catarrhalis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct

15 expression of BASB034 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB034 polynucleotide and/or polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease,

20 whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

25 A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB034 polynucleotide and/or polypeptide encoded therefrom, wherein the

composition comprises a recombinant BASB034 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB034 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically
5 or prophylactically and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

A BASB034 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of
10 stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and
15 facilitates production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

20 Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

25 Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Moraxella catarrhalis*. Such experiments will be particularly useful for identifying protein epitopes able to

provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial
5 infection, particularly *Moraxella catarrhalis* infection, in mammals, particularly humans.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and
10 polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostatic compounds and solutes which render the formulation isotonic with
15 the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

20

The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

25 An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (*Mosmann, T.R. and Coffman, R.L. (1989) TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology, 7, p145-173*). Traditionally, TH1-type responses are associated with the production of the INF- γ and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2-type responses are associated with the secretion of IL-4, IL-5, IL-6 and IL-13.

It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2-type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes

direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes *in vitro* after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

- 5 Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell populations to produce high levels of TH1-type cytokines when re-stimulated with antigen *in vitro*, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

- 10 Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

- 3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated
15 monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem, Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

- Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a
20 0.22micron membrane (European Patent number 0 689 454).
3D-MPL will be present in the range of 10µg - 100µg preferably 25-50µg per dose wherein the antigen will typically be present in a range 2-50µg per dose.

- Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction
25 derived from the bark of Quillaja Saponaria Molina. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with an carrier.

The method of production of QS21 is disclosed in US patent No. 5,057,540.

Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; preferably 1:5 to 5 : 1 and often substantially 1 : 1. The preferred range for optimal synergy is 2.5 : 1 to 1 : 1 3D-MPL: QS21.

Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1µg - 200µg, such as 10-100µg, preferably 10µg - 50µg per dose.

Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene: alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85 may also be present at a level of 1%. In some cases it may be advantageous that the
5 vaccines of the present invention will further contain a stabiliser.

Non-toxic oil in water emulsions preferably contain a non-toxic oil, e.g. squalane or squalene, an emulsifier, e.g. Tween 80, in an aqueous carrier. The aqueous carrier may be, for example, phosphate buffered saline.
10

A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil in water emulsion is described in WO 95/17210.

The present invention also provides a polyvalent vaccine composition comprising a
15 vaccine formulation of the invention in combination with other antigens, in particular antigens useful for treating cancers, autoimmune diseases and related conditions. Such a polyvalent vaccine composition may include a TH-1 inducing adjuvant as hereinbefore described.

20 While the invention has been described with reference to certain BASB034 polypeptides and polynucleotides, it is to be understood that this covers fragments of the naturally occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant polypeptides or polynucleotides.

25

Compositions, kits and administration

In a further aspect of the invention there are provided compositions comprising a BASB034 polynucleotide and/or a BASB034 polypeptide for administration to a cell or to a multicellular organism.

The invention also relates to compositions comprising a polynucleotide and/or a polypeptides discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or
5 sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to an individual. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol,
10 ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

15 Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous,
20 intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

25 In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such as the soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically

acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

- 5 Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

- The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by
10 intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral
15 administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.

- For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1
20 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

- 25 The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

15

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination.

nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

5 A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

10 A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

15 All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference
20 herein in its entirety in the manner described above for publications and references.

DEFINITIONS

25 "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known

- methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heine, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are
- 10 codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990), and FASTA(Pearson and Lipman *Proc. Natl. Acad. Sci. USA* 85: 2444-2448
- 15 (1988). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman algorithm may also be used to determine identity.
- 20 Parameters for polypeptide sequence comparison include the following:
Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)
Comparison matrix: BLOSSUM62 from Henikoff and Henikoff,
Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)
Gap Penalty: 8
- 25 Gap Length Penalty: 2
- A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

Algorithm: Needleman and Wunsch, J. Mol Biol. 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

5 Gap Length Penalty: 3

Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

10 A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

(1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide
 15 sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference
 20 nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that
 25 product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y).$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded
 5 down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

- 10 By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion,
 15 substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent
 20 identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

25

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \cdot is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion.

substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one
 5 or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$10 \quad n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and
 15 y is rounded down to the nearest integer prior to subtracting it from x_a .

"Individual(s)," when used herein with reference to an organism, means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

20 "Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural
 25 state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism. which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA including single and double-stranded regions.

5 "Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid
10 sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions,
15 identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally
20 occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Disease(s)" means any disease caused by or related to infection by a bacteria, including, for example, otitis media in infants and children, pneumonia in elderlies, sinusitis,
25 nosocomial infections and invasive diseases, chronic otitis media with hearing loss, fluid accumulation in the middle ear, auditive nerve damage, delayed speech learning, infection of the upper respiratory tract and inflammation of the middle ear.

EXAMPLES:

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1:**Discovery and confirmatory DNA sequencing of the BASB034 gene from *Moraxella catarrhalis* strain ATCC 43617.**

10

The BASB034 gene was first discovered in the Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Moraxella catarrhalis* strain ATCC 43617 (also referred to as strain MC2931). The translation of the BASB034 polynucleotide sequence showed significant similarity (42 % identity in a 214 amino acids overlap) to a *Klebsiella pneumoniae* outer membrane phospholipase A protein.

The sequence of the BASB034 gene was further confirmed experimentally. For this purpose, genomic DNA was extracted from 10^{10} cells of the *M. catarrhalis* cells (strain ATCC 43617) using the QIAGEN genomic DNA extraction kit (Qiagen GmbH), and 1 µg of this material was submitted to Polymerase Chain Reaction DNA amplification using primers E481124 (5'- GAT TTA AGA GTA TGT TAT GAT G -3') [SEQIDNO:9] and E481125 (5'-GTA TGG GTT GAT CAA ATA CAG-3') [SEQIDNO:10]. This PCR product was purified on a Biorobot 9600 (Qiagen GmbH) apparatus and subjected to DNA sequencing using the Big Dye Cycle Sequencing kit (Perkin-Elmer) and an ABI 377/PRISM DNA sequencer. DNA sequencing was performed on both strands with a redundancy of 2 and the full-length sequence was assembled using the Sequencher™ software (Applied Biosystems). The resulting DNA sequence turned out to be 100% identical to SEQ ID NO:1.

Example 2:

Variability analysis of the BASB034 gene among several *Moraxella catarrhalis* strains.

5

2A: Restriction Fragment Length Analysis (RFLP).

Genomic DNA was extracted from 16 *M. catarrhalis* strains (presented in Table 1) as described below. *M. catarrhalis* was streaked for single colonies on BHI agar plates and grown overnight at 37 °C. Three or four single colonies were picked and used to
10 inoculate a ~1.5 ml BHI (Brain-heart infusion) broth seed culture which was grown overnight in a shaking incubator, ~300 rpm, at 37 °C. A 500ml erlenmeyer flask containing ~150 ml of BHI broth was inoculated with the seed culture and grown for ~12-16 hours at 37 °C in a shaking incubator, ~175 rpm, to generate cell mass for DNA isolation. Cells were collected by centrifugation in a Sorvall GSA rotor at ~2000 X g
15 for 15 minutes at room temperature. The supernatant was removed and the cell pellet suspended in ~5.0 ml of sterile water. An equal volume of lysis buffer (200 mM NaCl, 20 mM EDTA, 40 mM Tris-Hcl, pH 8.0, 0.5% (w/v) SDS, 0.5% (v/v) 2-mercaptoethanol, and 250 µg/ml of proteinase K) was added and the cells suspended by gentle agitation and trituration. The cell suspension was then incubated ~12 hours at
20 50°C to lyse the bacteria and liberate chromosomal DNA. Proteinaceous material was precipitated by the addition of 5.0 ml of saturated NaCl (~6.0 M, in sterile water) and centrifugation at ~5,500xg in a Sorvall SS34 rotor at room temperature. Chromosomal DNA was precipitated from the cleared supernatant by the addition of two volumes of 100 % ethanol. Aggregated DNA was collected and washed using gentle agitation in a
25 small volume of a 70 % ethanol solution. Purified chromosomal DNA was suspended in sterile water and allowed to dissolve/ disburse overnight at 4 °C by gentle rocking. The concentration of dissolved DNA was determined spectrophotometrically at 260 nm using an extinction coefficient of 1.0 O.D. unit ~50 µg/ml.

This material was next submitted to PCR amplification using the MC-Pla-BamF (5'-AAG GGC CCA ATT ACG CAG AGG GGA TCC CAA GCT GTA CCA AAT CCT GTG GCA TTT GTT G -3') [SEQ ID NO:11] and MC-Pla-SalRC (5'-AAG GGC CCA ATT ACG CAG AGG GTC GAC TTA TTA TAG ACC CAT CCA GTC GTT AAG CAT AAG -3') [SEQ ID NO:12] oligonucleotides. The corresponding BASB034 gene amplicons were then subjected independently to hydrolysis using restriction enzymes (*HphI*, *AluI*, *Rsa I*, *EcoRV*, *Sau3AI*) and restriction products were separated by agarose or polyacrylamide gel electrophoresis using standard molecular biology procedures as described in "Molecular Cloning, a Laboratory Manual, Second Edition, Eds: Sambrook, Fritsch & Maniatis, Cold Spring Harbor press 1989". The photographs of the resulting electrophoresis gels are displayed in Figure 1. For each strain, RFLP patterns corresponding to the 6 restriction enzymes were scored and combined. Groups of strains sharing identical combination of RFLP patterns were then defined. Using this methodology, the strains tested in this study fell into 3 genomic groups (Group 1: Mc2931, Mc 2904, Mc 2905, Mc2907, Mc2909, Mc2910, Mc2911, Mc2912, Mc2926, Mc2931, Mc2956, Mc2969, Mc2975; Group 2: Mc2906, Mc2913; Group 3: Mc2908. (Mc2960 failed to be amplified and consequently was not classified). These data support that the *Moraxella catarrhalis* population used in this study displays limited nucleotide sequence diversity for the BASB034 gene.

2B: DNA sequencing in other strains.

Using the experimental procedure described in Example 1, the sequence of the BASB034 gene was also determined for three additional *Moraxella catarrhalis* strains. The nucleotide sequences of the BASB034 gene of the strains Mc2908, Mc2913 and Mc2969, representative of the three genomic groups identified previously, are shown in SEQ ID NO:3, 5 and 7, respectively. These nucleotide sequences were translated into amino acid sequences, which are shown in SEQ ID NO:4, 6 and 8, respectively. Using the MegAlign program from the DNASTAR Lasergene package, a multiple alignment of the nucleotide sequences of SEQ ID NO:1, 3, 5 and 7 was performed, and is

displayed in Figure 2. A pairwise comparison of identities is summarized in Table 2, showing that the four BASB034 nucleotide gene sequences are all similar at an identity level greater than 98 %. Using the same program, a multiple alignment of the protein sequences of SEQ ID NO:2, 4, 6 and 8 was performed, and is displayed in Figure 3. A pairwise comparison of identities is summarized in Table 3, showing that the four BASB034 protein sequences are all similar at an identity level greater than 98 %. Taken together, these data indicate very strong sequence conservation of the BASB034 gene among *Moraxella catarrhalis* strains.

10 **Table 1: Features of the *Moraxella catarrhalis* strains used in this study**

Strain	Isolated in:	from:
Mc2904	USA	Tympanocentesis
Mc2905	USA	Tympanocentesis
Mc2906	USA	Tympanocentesis
Mc2907	USA	Tympanocentesis
Mc2908	USA	Acute otitis Tympanocentesis
Mc2909	USA	Tympanocentesis
Mc2910	USA	Tympanocentesis
Mc2911	USA	Acute otitis Tympanocentesis
Mc2912	USA	Acute otitis Tympanocentesis
Mc2913	USA	Acute otitis Tympanocentesis
Mc2926	USA	Tympanocentesis
Mc2931 /ATCC 43617	USA	Transtracheal aspirate
Mc2956	Finland	Middle ear fluid
Mc2960	Finland	Middle ear fluid
Mc2969	Norway	Nasopharynx (Pharyngitis-Rhinitis)
Mc2975	Norway	Nasopharynx (Rhinitis)

Table 2: Pairwise identities of the BASB034 polynucleotide sequences (in %)

	SeqID No:3	SeqID No:5	SeqID No:7
SeqID No:1	98.7	99.2	99.7
SeqID No:3		99.3	98.7
SeqID No:5			99.1

5 Table 3. Pairwise identities of the BASB034 polypeptide sequences (in %)

	SeqID No:4	SeqID No:6	SeqID No:8
SeqID No:2	98.6	99.3	99.5
SeqID No:4		98.9	99.1
SeqID No:6			99.3

Example 3: Construction of Plasmid to Express Recombinant BASB034

5

A: Cloning of BASB034.

The *Bam*HI and *Sal*I restriction sites engineered into the forward ([SEQ ID NO:11]) and reverse ([SEQ ID NO:12]) amplification primers, respectively, permitted directional cloning of a BASB034 PCR product into the commercially available *E.coli* expression plasmid pQE30 (QiaGen, ampicillin resistant) such that a mature BASB034 protein could be expressed as a fusion protein containing a (His)₆ affinity chromatography tag at the N-terminus. The BASB034 PCR product was purified from the amplification reaction using silica gel-based spin columns (QiaGen) according to the manufacturers instructions. To produce the required *Bam*HI and *Sal*I termini necessary for cloning, 10 purified PCR product was sequentially digested to completion with BamHI and SalI 15 restriction enzymes as recommended by the manufacturer (Life Technologies). Following the first restriction digestion, the PCR product was purified via spin column as above to remove salts and eluted in sterile water prior to the second enzyme digestion. The digested DNA fragment was again purified using silica gel-based spin 20 columns prior to ligation with the pQE30 plasmid.

B: Production of Expression Vector.

To prepare the expression plasmid pQE30 for ligation, it was similarly digested to completion with both *Bam*HI and *Sal*I and then treated with calf intestinal phosphatase 25 (CIP, ~0.02 units / pmole of 5' end, Life Technologies) as directed by the manufacturer to prevent self ligation. An approximately 5-fold molar excess of the digested fragment to the prepared vector was used to program the ligation reaction. A standard ~20 µl ligation reaction (~16°C, ~16 hours), using methods well known in the art, was performed using T4 DNA ligase (~2.0 units / reaction, Life Technologies). An aliquot 30 of the ligation (~5 µl) was used to transform electro-competent M15(pREP4) cells

according to methods well known in the art. Following a ~2-3 hour outgrowth period at 37°C in ~1.0 ml of LB broth, transformed cells were plated on LB agar plates containing kanamycin (50 µg/ml) and ampicillin (100 µg/ml). Both antibiotics were included in the selection media to ensure that all transformed cells carried both the

5 pREP4 plasmid (KnR), which carries the *lacIq* gene necessary for the repression of expression for IPTG-inducible expression of proteins on pQE30, and the pQE30-BASB034 plasmid (ApR). Plates were incubated overnight at 37°C for ~16 hours. Individual KnR / ApR colonies were picked with sterile toothpicks and used to "patch" inoculate fresh LB KnR / ApR plates as well as a ~1.0 ml LB KnR / ApR broth culture.

10 Both the patch plates and the broth culture were incubated overnight at 37°C in either a standard incubator (plates) or a shaking water bath.

A whole cell-based PCR analysis was employed to verify that transformants contained the BASB034 DNA insert. Here, the ~1.0 ml overnight LB Kn / Ap broth culture was

15 transferred to a 1.5 ml polypropylene tube and the cells collected by centrifugation in a Beckman microcentrifuge (~3 min., room temperature, ~12,000 X g). The cell pellet was suspended in ~200µl of sterile water and a ~10µl aliquot used to program a ~50µl final volume PCR reaction containing both BASB034 forward and reverse amplification primers. Final concentrations of the PCR reaction components were essentially the

20 same as those specified in example 2 except ~5.0 units of *Taq* polymerase was used. The initial 95°C denaturation step was increased to 3 minutes to ensure thermal disruption of the bacterial cells and liberation of plasmid DNA. An ABI Model 9700 thermal cycler and a 32 cycle, three-step thermal amplification profile, i.e. 95°C, 45sec; 55-58°C, 45sec, 72°C, 1min., were used to amplify the BASB034 PCR fragment from

25 the lysed transformant samples. Following thermal amplification, a ~20µl aliquot of the reaction was analyzed by agarose gel electrophoresis (0.8 % agarose in a Tris-acetate-EDTA (TAE) buffer). DNA fragments were visualized by UV illumination after gel electrophoresis and ethidium bromide staining. A DNA molecular size standard (1 Kb ladder, Life Technologies) was electrophoresed in parallel with the test samples and was

used to estimate the size of the PCR products. Transformants that produced the expected PCR product were identified as strains containing a BASB034 expression construct. Expression plasmid containing strains were then analyzed for the inducible expression of recombinant BASB034.

5

C: Expression Analysis of PCR-Positive Transformants.

For each PCR-positive transformant identified above, ~5.0 ml of LB broth containing kanamycin (50 µg/ml) and ampicillin (100 µg/ml) was inoculated with cells from the patch plate and grown overnight at 37 °C with shaking (~250 rpm). An aliquot of the
10 overnight seed culture (~1.0 ml) was inoculated into a 125 ml erlenmeyer flask containing ~25 ml of LB Kn / Ap broth and grown at 37 °C with shaking (~250 rpm) until the culture turbidity reached O.D.600 of ~0.5, i.e. mid-log phase (usually about 1.5 - 2.0 hours). At this time approximately half of the culture (~12.5 ml) was transferred to a second 125 ml flask and expression of recombinant BASB034 protein induced by
15 the addition of IPTG (1.0 M stock prepared in sterile water, Sigma) to a final concentration of 1.0 mM. Incubation of both the IPTG-induced and non-induced cultures continued for an additional ~4 hours at 37 °C with shaking. Samples (~1.0 ml) of both induced and non-induced cultures were removed after the induction period and the cells collected by centrifugation in a microcentrifuge at room temperature for ~3
20 minutes. Individual cell pellets were suspended in ~50µl of sterile water, then mixed with an equal volume of 2X Laemmli SDS-PAGE sample buffer containing 2-mercaptoethanol, and placed in boiling water bath for ~3 min to denature protein. Equal volumes (~15µl) of both the crude IPTG-induced and the non-induced cell lysates were loaded onto duplicate 12% Tris/glycine polyacrylamide gel (1 mm thick Mini-gels,
25 Novex). The induced and non-induced lysate samples were electrophoresed together with prestained molecular weight markers (SeeBlue, Novex) under conventional conditions using a standard SDS/Tris/glycine running buffer (BioRad). Following electrophoresis, one gel was stained with commassie brilliant blue R250 (BioRad) and then destained to visualize novel BASB034 IPTG-inducible protein(s). The second gel

was electroblotted onto a PVDF membrane (0.45 micron pore size, Novex) for ~2 hrs at 4 °C using a BioRad Mini-Protean II blotting apparatus and Towbin's methanol (20 %) transfer buffer. Blocking of the membrane and antibody incubations were performed according to methods well known in the art. A monoclonal anti-RGS (His)₃ antibody, followed by a second rabbit anti-mouse antibody conjugated to HRP (QiaGen), was used to confirm the expression and identity of the BASB034 recombinant protein. Visualization of the anti-His antibody reactive pattern was achieved using either an ABT insoluble substrate or using Hyperfilm with the Amersham ECL chemiluminescence system.

10

D: Sequence Confirmation.

To further verify that the IPTG-inducible recombinant BASB034 protein being expressed is in the correct open reading frame and not a spurious molecule arising from a cloning artifact (i.e. a frame-shift), the DNA sequence of the cloned insert was determined. The DNA sequence for the *M.catarrhalis* BASB034 gene was obtained from one strand using conventional asymmetric PCR cycle sequencing methodologies (ABI Prism Dye-Terminator Cycle Sequencing, Perkin-Elmer). Sequencing reactions were programmed with undigested expression plasmid DNA (~0.5µg/rxn) as a template and appropriate pQE30 vector-specific and ORF-specific sequencing primers (~3.5 pmol/rxn). In addition to the template and sequencing primer, each sequencing reaction (~20µl) contained the four different dNTPs (i.e. A,G,C, and T) and the four corresponding ddNTPs (i.e. ddA, ddG, ddC, and ddT) terminator nucleotides; with each terminator being conjugated to one of the four fluorescent dyes, Joe, Tam, Rox, or Fam. Single strand sequencing elongation products were terminated at random positions along the template by the incorporation of the dye-labelled ddNTP terminators. Fluorescent dye-labelled termination products were purified using microcentrifuge size-exclusion chromatography columns (Princeton Genetics), dried under vacuum, suspended in a Template Resuspension Buffer (Perkin-Elmer) for capillary electrophoresis or deionized formamide for PAGE. denatured at 95°C for ~5 min. and

analyzed by high resolution capillary electrophoresis (ABI 310 Automated DNA Sequenator, Perkin-Elmer) or high resolution PAGE (ABI 377 Automated DNA Sequenator) as recommended by the manufacturer. DNA sequence data produced from individual reactions were collected and the relative fluorescent peak intensities analyzed automatically on a PowerMAC computer using ABI Sequence Analysis Software (Perkin-Elmer). Individually autoanalyzed DNA sequences were edited manually for accuracy before being merged into a consensus single strand sequence "string" using AutoAssembler software (Perkin-Elmer). Sequencing determined that the expression plasmid contained the correct sequence in the correct open reading frame.

10

Example 4: Production of Recombinant BASB034

Bacterial strain

15 A recombinant expression strain of *E. coli* M15 (pREP4) containing a plasmid (pQE30) encoding BASB034 from *M. catarrhalis*. was used to produce cell mass for purification of recombinant protein. The expression strain was cultivated on LB agar plates containing 50µg/ml kanamycin ("Kn") and 100µg/ml ampicillin ("Ap") to ensure both the pREP4 lacIq control plasmid and the pQE30-BASB034 expression construct were
20 both maintained. For cryopreservation at -80 °C, the strain was propagated in LB broth containing the same concentration of antibiotics then mixed with an equal volume of LB broth containing 30% (w/v) glycerol.

Media

25 The fermentation medium used for the production of recombinant protein consisted of 2X YT broth (Difco) containing 50µg/ml Kn and 100µg/ml Ap. Antifoam was added to medium for the fermentor at 0.25 ml/L (Antifoam 204, Sigma). To induce expression of the BASB034 recombinant protein, IPTG (Isopropyl β-D-Thiogalactopyranoside) was added to the fermentor (1 mM, final).

30

Fermentation

A 500-ml erlenmeyer seed flask, containing 50ml working volume, was inoculated with 0.3 ml of rapidly thawed frozen culture, or several colonies from a selective agar plate culture, and incubated for approximately 12 hours at $37 \pm 1^\circ\text{C}$ on a shaking platform at 150rpm (Innova 2100, New Brunswick Scientific). This seed culture was then used to inoculate a 5-L working volume fermentor containing 2X YT broth and both Kn and Ap antibiotics. The fermentor (Bioflo 3000, New Brunswick Scientific) was operated at $37 \pm 1^\circ\text{C}$, 0.2 - 0.4 VVM air sparge, 250 rpm in Rushton impellers. The pH was not controlled in either the flask seed culture or the fermentor. During fermentation, the pH ranged 6.5 to 7.3 in the fermentor. IPTG (1.0 M stock, prepared in sterile water) was added to the fermentor when the culture reached mid-log of growth (~ 0.7 O.D.600 units). Cells were induced for 2 - 4 hours then harvested by centrifugation using either a 28RS Heraeus (Sepatech) or RC5C superspeed centrifuge (Sorvall Instruments). Cell paste was stored at -20°C until processed.

Purification

Chemicals and Materials

Imidazole, guanidine hydrochloride, Tris (hydroxymethyl), and EDTA (ethylene-diamine tetraacetic acid) biotechnology grade or better were all obtained from Ameresco Chemical, Solon, Ohio. Triton X-100 (t-Octylphenoxypolyethoxy-ethanol), sodium phosphate, monobasic, and Urea were reagent grade or better and obtained from Sigma Chemical Company, St. Louis, Missouri. Glacial acetic acid and hydrochloric acid were obtained from Mallincrodt Baker Inc., Phillipsburg, New Jersey. Methanol was obtained from Fisher Scientific, Fairlawn, New Jersey. Pefabloc®SC (4-(2-Aminoethyl)-benzenesulfonylfluoride), Complete protease inhibitor cocktail tablets, and PMSF (phenylmethyl-sulfonylfluoride) were obtained from Roche Diagnostics Corporation, Indianapolis, Indiana. Bestatin, Pepstatin A, and E-64 protease inhibitor were obtained from Calbiochem, LaJolla, California. Dulbecco's Phosphate Buffered Saline (1x PBS) was obtained from Quality Biological, Inc., Gaithersburg, Maryland.

Dulbecco's Phosphate Buffered Saline (10x PBS) was obtained from BioWhittaker, Walkersville, Maryland. Penta-His Antibody, BSA free was obtained from QiaGen, Valencia, California. Peroxidase-conjugated AffiniPure Goat Anti-mouse IgG was obtained from Jackson Immuno Research, West Grove, Penn. AEC single solution was
5 obtained from Zymed, South San Francisco, California. All other chemicals were reagent grade or better.

Ni-NTA Superflow resin was obtained from QiaGen Inc., Valencia, California. Precast Tris-Glycine 4-20% and 10-20% polyacrylamide gels, all running buffers and solutions, SeeBlue Pre-Stained Standards, MultiMark Multi-Colored Standards and PVDF transfer
10 membranes were obtained from Novex, San Diego, California. SDS-PAGE Silver Stain kits were obtained from Daiichi Pure Chemicals Company Limited, Tokyo, Japan. Coomassie Stain Solution was obtained from Bio-Rad Laboratories, Hercules, California. Acrodisc® PF 0.2 µm syringe filters were obtained from Pall Gelman Sciences, Ann Arbor, Michigan. GD/X 25mm disposable syringe filters were obtained
15 from Whatman Inc., Clifton, New Jersey. Dialysis tubing 8,000 MWCO was obtained from BioDesign Inc. Od New York, Carmal New York. BCA Protein Assay Reagents and Snake Skin dialysis tubing 3,500 MWCO were obtained from Pierce Chemical Co., Rockford, Illinois.

20 Extraction Protocol

Cell paste was thawed at room temperature for 30 to 60 minutes. Five to six grams of material was weighed out into a 50 ml disposable centrifuge tube. To this five
mls/gram of Guanidine hydrochloride (Gu-HCl) buffer was added (6 M Guanidine hydrochloride, 100 mM Sodium phosphate, monobasic, 10 mM Tris and 0.05 % Triton
25 X-100, pH 8.0). Cell paste was resuspended using a PRO300D proscientific homogenizer, at 3/4 power for one minute. The extraction mixture was then placed at room temperature with gentle agitation for 60 to 90 minutes. After 60 to 90 minutes the extraction mixture was centrifuged at 15,800 x g for 15 minutes (Sorvall RC5C

centrifuge, 11,500 rpm). The supernatant (S1) was decanted and saved for additional purification. The pellet (P1) was saved for analysis.

Binding of BASB034 to Nickel-NTA Resin

- 5 To the S1 three to four mls of Ni-NTA resin is added. This is then placed at room temperature with gentle agitation for one hour. After one hour the S1/Ni-NTA is packed into an XK16 Pharmacia column. The column is then washed with 1 M Gu-HCl buffer (1 M Guanidine hydrochloride, 100mM Sodium phosphate, monobasic, 10 mM Tris and 0.05% Triton X-100, pH 8.0). This is then followed by a wash with phosphate
- 10 buffer (100mM Sodium phosphate, monobasic, 10 mM Tris and 0.05% Triton X-100, pH 6.3). The protein is then eluted from the column with a 250 mM imidazole buffer (250 mM imidazole, 100mM Sodium phosphate, monobasic, 10 mM Tris and 0.05% Triton X-100, pH 5.9).

15 Final Formulation

BASB034 was formulated by dialysis overnight against, three changes of 0.1 % Triton X-100 and 1x PBS, pH 7.4, to remove residual Gu-HCl and imidazole. The purified protein was characterized and used to produced antibodies as described below.

20 Biochemical Characterizations

SDS-PAGE and Western Blot Analysis

- The recombinant purified protein was resolved on 4-20 % polyacrylamide gels and electrophoretically transferred to PVDF membranes at 100 V for 1 hour as previously described (Thebaine et al. 1979, Proc. Natl. Acad. Sci. USA 76:4350-4354). The
- 25 PVDF membranes were then pretreated with 25 ml of Dulbecco's phosphate buffered saline containing 5 % non-fat dry milk. All subsequent incubations were carried out using this pretreatment buffer.

- PVDF membranes were incubated with 25 ml of a 1:500 dilution of preimmune serum or rabbit anti-His immune serum for 1 hour at room temperature. PVDF membranes were then washed twice with wash buffer (20 mM Tris buffer, pH 7.5, containing 150 mM sodium chloride and 0.05 % Tween-20). PVDF membranes were incubated with
- 5 25 ml of a 1:5000 dilution of peroxidase-labeled goat anti-rabbit IgG (Jackson ImmunoResearch Laboratories, West Grove, PA) for 30 minutes at room temperature. PVDF membranes were then washed 4 times with wash buffer, and were developed with 3-amino-9-ethylcarbazole and urea peroxide as supplied by Zymed (San Francisco, CA) for 10 minutes each.
- 10 The results of an SDS-PAGE (Figure 4) show a protein about 60 kDa that is reactive to an anti-RGS(His) antibody by western blots (Figure 5) of the SDS-PAGE.

Protein Sequencing

- Amino terminal amino acid sequencing of the purified protein was performed to
- 15 confirm the production of the correct recombinant protein using well defined chemical protocols on Hewlett-Packard model G1000A sequencer with a model 1090 LC and a Hewlett-Packard model 241 sequencer with a model 1100 LC.

Example 5: Production of Antisera to Recombinant BASB034

- 20 Polyvalent antisera directed against the BASB034 protein were generated by vaccinating two rabbits with the purified recombinant BASB034 protein. Each animal is given a total of three immunizations intramuscularly (i.m.) of about 20µg BASB034 protein per injection (beginning with complete Freund's adjuvant and followed with incomplete Freund's adjuvant) at approximately 21 day intervals. Animals were bled
- 25 prior to the first immunization ("pre-bleed") and on days 35 and 57.

Anti-BASB034 protein titres were measured by an ELISA using purified recombinant BASB034 protein (0.5 µg/well). The titre is defined as the highest dilution equal or

greater than 0.1 as calculated with the following equation: average OD of two test samples of antisera - the average OD of two test samples of buffer. The titres after three immunizations were around 1000000.

The antisera were used as the first antibody to identify the protein in a western blot as described in example 4 above. The western-blot shows the presence of anti-BASB034 antibody in the sera of immunized animals (Figure 6).

Example 6 : Analysis of the non-coding flanking regions of the BASB034 gene, and its exploitation for modulated BASB034 gene expression.

The non-coding flanking regions of the BASB034 gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the shine-dalgarno sequence, regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences.

This sequence information allows the modulation of the natural expression of gene BASB034. The upregulation of the gene expression may be accomplished by altering the promoter, the shine-dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modifications. Alternatively, by changing phase variation sequences, the expression of the gene can be put under phase variation control, or may be uncoupled from this regulation. In another approach, the expression of the gene can be put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation include, but are not limited to, induction by temperature shift, addition of inductor substrates

like selected carbohydrates or their derivatives, trace elements, vitamins, co-factors, metal ions, etc.

Such modifications as described above can be introduced by several different means.

The modification of sequences involved in gene expression can be done *in vivo* by

- 5 random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating the region of interest and modifying it by random mutagenesis, or site-directed mutagenesis, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge
- 10 of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the wild-type regulatory sequences.
- 15 These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome. A non-exhaustive list of preferred promoters that could be used for up-regulation of gene expression includes the promoter *porA*, *porB*, *lbpB*, *tbpB*, *p110*, *1st*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*.
- In one example, the expression of the gene can be modulated by exchanging its
- 20 promoter with a stronger promoter (through isolating the upstream sequence of the gene, *in vitro* modification of this sequence, and reintroduction into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium.
- In other examples, the described approaches can be used to generate recombinant bacterial
- 25 strains with improved characteristics for vaccine applications. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the immune response, strains with modulated expression of immunodominant proteins, strains with modulated shedding of outer-membrane vesicles.

A region directly upstream of the BASB034 gene is given in the sequence of SEQ ID NO:13. This sequence is a further aspect of the invention.

SEQUENCE INFORMATION

BASB034 Polynucleotide and Polypeptide Sequences

SEQ ID NO:1

5 *Moraxella catarrhalis* BASB034 polynucleotide sequence from strain ATCC 43617

ATGAAAGTTTCACTGTCTACATTGACTTTATCTATTTTGTGCTATCCTAGCCATTGAGCAAGCACAGCTGT
 ACCAAATCCTGTGGCATTGTTGACGAAGTACGCAGTGAAAATGATCTTGGGCAAGACAATGAATTACCCATTGATGTCC
 AAAGTGCACACAATCAGCGTCTACTGATACGGCTAATCCTTTAGACGAACATGAACCAGAGCTTTATACGACAGCTTTA
 10 GAAAATAAAACCATGCTGATTAACTGCTCAGCACTTAATCAAGATATCATGCGTTTGGCGTGCTATGACACTTTGGTGCA
 TGGTGAGACGCCAGCGTAATTAACCAAGCGTCCATTGCGCTTGATGAAACAATTTGGCAGACCATCAAAGGCAAAC
 CCCAGGTTATCTATCAAGAAACGACAGATCCGATTTTTTAATGGGTAATGAAAAAGGCATGCTGACCAAAAAAGATGCC
 AAACAGCTTGAATATGCAGCCAAACAGTTTACACCACTGAGCTTATCATTTGATTAGACCGAAATAATACACCACTTTG
 GTCATCACGACCACACAATCCGATGTATGTATTGCCATATTTATGCACGGTAAGCCTAATCGAAGCCCAAATACGCCCA
 15 GTCATGAAGCAAAACAATTTACCCCAAATGAATTTGCTGCTCCCGAGCTAAAATTTGAGTTTCTGTTAAGGTTAAAGCT
 GCTGAGGATTTATGGGGGACGGATTGAGTTTATGGTTTGGATATACACAGCAATCGCACTGGCAGATTTTTAATGGAAA
 AAATCTCGTCTTTTAGAGTACATGACTACCAGCCAGAGATTTTCTTAACCTCAACCTGTATACTCAGACTTACCATGGG
 ATGGCAAAGTCCGATGATTGGCATGGGTGCGGTACATCATTCCAATGGTGAAAGTGCCAACTGTCTCGCTCATGGAAT
 CGTGCTTATTTGATGGCAGGCATGGAATGAAAAACCTGACTGTATGCCACGCATTTGGGGCGTATCTTTAAAGAGGG
 20 TAGTGGCAGCCAGCCAGATGATAATCCTGATATCTGGACTATTATGGTTATGGTGATGTGCGTTTTTATATCAACTAG
 AAAATAAAAGTAATATTTGAGGTACGGTACGCTATAATCCACGCTCAGGCAAAGGTGCGTTGCAACTTGACTATGTCTAT
 CCGCTTGGTAAGGAATTAGTGGCTATTTTCAAATATTTCAAGGCTATGGGCAGTCTTTGATTGATTATAATCATGAGGC
 GACAAGCTTTGGCGTCGGACTTATGCTTAACGACTGGATGGGTCTATAA

25 SEQ ID NO:2

Moraxella catarrhalis BASB034 polypeptide sequence deduced from the polynucleotide of
 Seq ID NO:1

30 MKVSLSTLTLSILSCFAILAIQQAQAVPNPFAFVDEVRSNDLQGDNELPIDVQSATQSASTDTANPLDEHEPELYTTAL
 ENKTMILINCSALNQDIMRLACYDTLVHGETPAVIKTKRSIRLDETIWQTIKGPQVIYQETTDPIFLMGNEKGMCLKKDA
 KQLEYAAKQFTPLSLSFDLDRNNTPLWSSRPHNPMYVLPIMHKGPNRSPNTPSHEAKQFTPNEFRAPELKFQVSVKVKKA
 AEDLWGTDSDLWFGYTQQSHWQIFNGKNSRPFVRVDYQPEIFLTQPVYSDLPWDGKVRMIGMGAHVHSHNGESAKLSRSWN
 RAYLMAGMEWKNLTVMPIWGRIFKEGSGSQPDNDPILDYYGYGDVRFYQLENKSNISGTVRYNPRSGKALQLDYVY
 35 PLGKGISGYFQIFQGYGQSLIDYNHEATSFGVGLMLNDWMGL

SEQ ID NO:3

Moraxella catarrhalis BASB034 polynucleotide sequence from strain Mc2908

ATGAAAGTTTCACTGTCTACATTGACTTTATCTATTTTCCCATGTTTTGCTATCCTAGCCATTGAGCAAGCACAGCTGT

ACCAAATCCTGTGGCATTGTGTGACGAAGTACGCAGTAAAAATGATCTTGGGCAAGACAATGAATTACTCATTGGTGTAC
 AAAGTGCAGACAAATCAGCGTCTACTGATACGGCTAATCCTTTAGACGAACATGAACCAGAGCTTTATACGACAGCTTTA
 GAAAAATAAACCATGCTGATTAACTGCTCAGCACTTAATCAAGATATCATGCGTTTGGCGTGCTATGACACTTTGGTGCA
 TGGTGAGACGCCAGCGTAATTAACAACCAAGCGTTCCATTGCGCTTGATGAAACAATTTGGCAGACCATCAAAGGCAAAC
 5 CCCAGGTTGTCTATCAAGAAACGACAGATCCGATTTTTTAAATGGGTAATGAAAAAGGCATGCTGACCAAAAAAGATGCC
 AAACAGCTTGAATATGCAGCCAAACAGTTTACACCACTGAGCTTATCATTTGATTAGACCGAAATAATACACCGCTTTG
 GTCATCAGGACACACAATCCGATGTATGTATTGCCATATTTATGCACGGTAAGCCTAATCGAAGCCCAATACGCCCA
 GTCATGAAGCAAGACAATTTACCCCAAATGAATTTGCGTGGCCCTGAATTAATAATTTCAAGTTTCTGTTAAGGTTAAAGCT
 GCTGAGGATTTATGGGGGACGGATTGAGATTATGGTTTGGGTATACACAGCAATCGCACTGGCAGATTTTAAATGGAAA
 10 AAATCTCGTCTTTTAGAGTACATGATTACCAGCCAGAGATTTTCTTAACCTCAACCTGTGTACTCAGACTTACCATGGG
 ATGGCAAAGTCCGATGATTGGCATGGGTGCGGTACATCATTCCAATGGTGAAAGTGCCAAACTGTCTCGCTCATGGAAT
 CGTGCTTATTTGATGGCAGGCATGGAATGGAATAAAGCTGACTGTATGCCACGCATTTGGGGGCGTATCTTTAAAGAGGG
 TAGTGGCAGCCAGCCAGATGACAATCCTGATATCTTGACTATTATGGTTATGGTGATGTGCGTTTTTATATCAACTAG
 AAAATAAAGTAATATTTAGGTACGGTACGTATAATCCACGCTCAGGCAAAGGTGCGTTGCACTTGACTATGTCTAT
 15 CCGCTTGGTAAGGAATTAGTGGCTATTTTCAAATATTTCAAGCTATGGGCAGTCTTTGATTGATTATAATCATGAGGC
 GACAAGCTTTGGCGTCGGACTTATGCTTAACGACTGGATGGGTCTATAA

SEQ ID NO:4

20 *Moraxella catarrhalis* BASB034 polypeptide sequence deduced from the polynucleotide of
 Seq ID NO:3

MKVSLSTLTLILPCFAILLAIQQAQAVPNPVAFFVDEVRSKNDLGQDNELLIGVQSATQSASTDTANPLDEHEPELYTTAL
 ENKTMILINCSALNQDIMRLACYDTLVHGETPAVIKTKRSIRLDETIWQTIKGKPVVYQETTDPIFLMGNEKGMLTKKDA
 KQLEYAAKQFTPLSLSFDLDRNNTPLWSSRPHPNMYVLPIMHGKPNRSPNTPSHEARQFTPNEFRAPELKFQVSVKVK
 25 AEDLWGTDSDLWFGYTTQSHWQIFNGKNSRPFVRVDYQPEIFLTQPVYSDLPWDGKVRMIGMAVHHSNGESAKLSRSWN
 RAYLMAGMEWKNLTVMPIWGRIFKEGSGSQPDNDPILDYGYGDVRFYQLNKSNI SGTVRYNPRSGKGALQLDYVY
 PLGKGISGYFQIFQGYGQSLIDYNHEATSFGVGLMLNDWMGL

SEQ ID NO:5

30 *Moraxella catarrhalis* BASB034 polynucleotide sequence from strain Mc2913

ATGAAAGTTTCACTGTCTACATTGACTTTATCTATTTTGTGATGTTTTGCTATCCTAGCCATTGAGCAAGCAAAGCTGT
 ACCAAATCCTGTGGCATTGTGTGACGAAGTACGCAGTAAAAATGATCTTGGGCAAGACAATGAATTACCCATTGATGTCC
 AAAGTGCAGACAAATCAGCGTCTACTGATACGGCTAATCCTTTAGACGAACATGAACCAGAGCTTTATACGACAGCTTTA
 35 GAAAAATAAACCATGCTGATTAACTGCTCAGCACTTAATCAAGATATCATGCGTTTGGCGTGCTATGACACTTTGGTGCA
 TGGTGAGACGCCAGCGTAATTAACAACCAAGCGTTCCATTGCGCTTGATGAAACAATTTGGCAGACCATCAAAGGCAAAC
 CCCAGGTTGTCTATCAAGAAACGACAGATCCGATTTTTTAAATGGGTAATGAAAAAGGCATGCTGACCAAAAAAGATGCC
 AAACAGCTTGAATATGCAGCCAAACAGTTTACACCACTGAGCTTATCATTTGATTAGACCGAAATAATACACCACTTTG
 GTCATCAGGACACACAATCCGATGTATGTATTGCCATATTTATGCACGGTAAGCCTAATCGAAGCCCAATACGCCCA
 40 GTCATGAAGCAAGACAATTTACCCCAAATGAATTTGCGTGGCCCTGAATTAATAATTTCAAGTTTCTGTTAAGGTTAAAGCT
 GCTGAGGATTTATGGGGGACGGATTGAGATTATGGTTTGGATATACACAGCAATCGCACTGGCAGATTTTAAATGGAAA

AAACTCTCGTCCTTTTAGAGTACATGATTACCAGCCAGAGATTTTCTTAACTCAACCTGTATACTCAGACTTACCATGGG
 ATGGCAAAGTCCGCATGATTGGCATGGGTGCGGTACATCATTCCAATGGTGAAAGTGCCAAACTGTCTCGCTCATGGAAT
 CGTGCTTATTTGATGGCAGGCATGGAATGAAAAACCTGACTGTATGCCACGCATTGGGGGCGTATCTTTAAAGAGGG
 TAGTGGCAGCCAGCCAGATGACAATCCTGATATCTTGGACTATTATGGTTATGGTGATGTGCGTTTTTATATCAACTAG
 5 AAAATAAAAGTAATATTTAGGTACGGTACGCTATAATCCACGCTCAGGCAAAGGTGCGTTGCAACTTGACTATGTCTAT
 CCGCTTGGTAAGGGAATTAGTGGCTATTTCAAATATTTCAAGGCTATGGGCAGTCTTTGATTGATTATAATCATGAGGC
 GACAAGCTTTGGCGTCGGACTTATGCTTAACGACTGGATGGGTCTATAA

SEQ ID NO:6

- 10 *Moraxella catarrhalis* BASB034 polypeptide sequence deduced from the polynucleotide of
 SeQ ID NO:5

MKVSLSLTLSILSCFAILAIQAKAVPNPVAFVDEVRENDLGQDNELPIDVQSATQSASTDTANPLDEHEPELYTTAL
 ENKTMILINCSALNQDIMRLACYDTLVHGETPAVIKTKRSIRLDETIWQTIKGPQVVYQETTDPIFLMGNEKGMLTKKDA
 15 KQLEYAAKQFTPLSLSFDLDRNNTPLWSSRPHNPMYVLPIMHKGPNRSPNTPSHEARQFTPNEFRAPELKFQVSVKVKA
 AEDLWGTDSDLNFGYTTQSHWQIFNGKNSRPFVRVDYQPEIFLTQPVYSDLPWDGKVRMIGMAVHHSNGESAKLSRSWN
 RAYLMAGMEWKNLTVMPRIWGRIFKEGSGSQPDNDPILDYGYGDVRFYQLENKSNISGTVRYNPRSGKALQLDYVY
 PLKGISGYFIQFGYQSLIDYNHEATSFGVGLMLNDWMGL

20 SEQ ID NO:7

Moraxella catarrhalis BASB034 polynucleotide sequence from strain Mc2969

ATGAAAGTTTCACTGTCTACATTGACTTTATCTATTTTGCCATGTTTGGCCATCCTAGCCATTGAGCAAGCACAAGCTGT
 ACCAAATCCTGTGGCATTGTTGACGAAGTACGCAGTGAAAATGATCTTGGGCAAGACAATGAATTACCCATTGATGTCC
 25 AAAGTGCAGACACAATCGGCGTCTACTGATACGGTAATCCTTTAGACGAACATGAACCAGAGCTTTATACGACAGCTTTA
 GAAAATAAAACCATGCTGATTAAGTCTCAGCACTTAATCAAGATATCATGCGTTTGGCGTGCTATGACACTTTGGTGCA
 TGGTGAGACGCCAGCGGTAATTAACCAAGCGTTCATTGCGCTTGATGAAACAATTTGGCAGACCATCAAAGGCCAAAC
 CCCAGGTTGTCTATCAAGAAACGACAGATCCGATTTTTTAAATGGGTAATGAAAAGGCATGCTGACCAAAAAGATGCC
 AAACAGCTTGAATATGCAGCCAAACAGTTTACACCACTGAGCTTATCATTGATTAGACCGAAATAATACACCACTTTG
 30 GTCATCAGCACCACACAATCCGATGTATGTATTGCCATATTTATGCACGGTAAGCCTAATCGAAGCCCAATACGCCCA
 GTCATGAAGCAAAACAATTTACCCCAAATGAATTTCTGTCTCCCGAGCTAAAATTTGAGTTTCTGTTAAGGTTAAAGCT
 GCTGAGGATTTATGGGGACGGATTGAGATTTATGGTTTGGATATACACAGCAATCGCACTGGCAGATTTTAAATGAAAA
 AAACCTCTCGTCCTTTTAGAGTACATGACTACCAGCCAGAGATTTTCTTAACTCAACCTGTATACTCAGACTTACCATGGG
 ATGGCAAAGTCCGCATGATTGGCATGGGTGCGGTACATCATTCCAATGGTGAAAGTGCCAAACTGTCTCGCTCATGGAAT
 35 CGTGCTTATTTGATGGCAGGCATGGAATGAAAAACCTGACTGTATGCCACGCATTGGGGGCGTATCTTTAAAGAGGG
 TAGTGGCAGCCAGCCAGATGATAATCCTGATATCTTGGACTATTATGGTTATGGTGATGTGCGTTTTTATATCAACTAG
 AAAATAAAAGTAATATTTAGGTACGGTACGCTATAATCCACGCTCAGGCAAAGGTGCGTTGCAACTTGACTATGTCTAT
 CCGCTTGGTAAGGGAATTAGTGGCTATTTCAAATATTTCAAGGCTATGGGCAGTCTTTGATTGATTATAATCATGAGGC
 GACAAGCTTTGGCGTCGGACTTATGCTTAACGACTGGATGGGTCTATAA

SEQ ID NO:8

Moraxella catarrhalis BASB034 polypeptide sequence deduced from the polynucleotide of
SeQ ID NO:7

5 MKVSLSTLTLSILPCFAILAIQQAQAVPNPVAFVDEVRENDLGQDNELPIDVQSATQSASTDTANPLDEHEPELYTTAL
ENKTMLINCSALNQDIMRLACYDTLVHGETPAVIKTKRSIRLDETIWQTIKGPQVYQETTDPIFLMGNEKGMLTKKDA
KQLEYAAKQFTPLSLSFDLDRNNTPLWSSRPHNPMYVLPIFMHGKPNRSPNTPSHEAKQFTPNEFRAPELKFQVSVKVKA
AEDLWGTDSDLWFGYTQQSHWQIFNGKNSRPFVVDYQPEIFLTQPVYSDLPWDGKVRMIGMGAVHHSNGESAKLSRSWN
10 RAYLMAGMEWKNTLVMPRIWGRIFKEGSGSQPDDNPDILDYGYGDVRFYQLENKSNISGTVRYNPRSGKALQLDYVY
PLGKGISGYFQIFQGYGQSLIDYNHEATSGVGLMLNDWMGL

SEQ ID NO:9

GAT TTA AGA GTA TGT TAT GAT G

15

SEQ ID NO:10

GTA TGG GTT GAT CAA ATA CAG

SEQ ID NO:11

20 AAG GGC CCA ATT ACG CAG AGG GGA TCC CAA GCT GTA CCA AAT CCT
GTG GCA TTT GTT G

SEQ ID NO:12

25 AAG GGC CCA ATT ACG CAG AGG GTC GAC TTA TTA TAG ACC CAT CCA
GTC GTT AAG CAT AAG

SEQ ID NO:13

30 ACTTGGCGAAAAATACCATTTATATCGATTGTGATGTTATACAGGCAGATGGCGGTACACGCACAGCCAGTATCAGTGGTG
CTGCGGTGGCACTTATTGATGCTTTAGAACACTTGACGCGTCGTAAAAAGCTTACCCAAGATCCGCTTTTGGGCTTGGTG
GCAGCGGTTTCTGTGGGTGTTAATCAAGGCCGTGATTGCTTGATTGGATTATGCTGAAGATTCAACTTGTGATACCGA
TTTAAATGTGGTCATGACGCAGGCAGGTGGGTTTATTGAGATTCAAGGCACAGCAGAAGAAAAGCCATTACTCGTGCTG
AAGCTAATGCGATGCTTGATTGGCAGAGCTGGGAATTGGGCAGATTATCGAAGCCCAAAGCAAGTATTAGGCTGGTGA
TATGCTAATCGTTGAAGATAATGGCGTGATCATCATTAAATGGACAAAGTAAAAGACCCATTATTTTGGTGGTCGATGA
35 TATTGCTGCTGCTGGGTGCTTGGTGGCAATCATTGTTTGGATTGCACCCGTTTTTTATGCAATCGGTGCGTTGGCTTTA
TTTGCACTTGTGGTATTGTGTTTAATATTCAAAGGCCAAAAGCCAAAACCTTGTCATATGTTTTACAAGGTCGCTTGAA
GATTACGTCCAAACGCTTTGAGATTATAACAAATCACTAACCTTATCAGCATCGGCAACAATATCTGCTAAAGATAACA
AAATGACAATTGTTGATCGGGGCATTGAATATCATTTTACAGGTTTTGCTGATGACCGTGAAATTAATATAGCCAAACAG
GTACTTTTGGGAAAAGTCAATCAAACCAATGCGGTGGCGGTAAACATTGGCTAAGTAGTTGTTGTGATACAGACAGGTGG
40 ATGGTCTTTAACTCCACCCACCTAACTTTTCTTTGTTGGATTAAAGAGTATGTTATGATGGGCAGGATTTTATTTTAA
GTCATCATTTAATGCAATCAGTTGTCCAGAGTAGCCGTTT

Deposited materials

5 A deposit containing a *Moraxella catarrhalis* Catlin strain has been deposited with the American Type Culture Collection (herein "ATCC") on June 21, 1997 and assigned deposit number 43617. The deposit was described as *Branhamella catarrhalis* (Frosch and Kolle) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *M. catarrhalis* isolate obtained from a transtracheal aspirate of a coal miner with chronic bronchitis. The deposit is described in Antimicrob. Agents Chemother. 21: 506-508 (1982).

10

The *Moraxella catarrhalis* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains a full length BASB034 gene.

15

A deposit of the vector pMC-PLA1 consisting of *Moraxella catarrhalis* DNA inserted in pQE30 has been deposited with the American Type Culture Collection (ATCC) on February 12 1999 and assigned deposit number 207099.

20 The sequence of the polynucleotides contained in the deposited strain, or in the deposited clone, as well as the amino acid sequence of any polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

25 The deposits of the deposited strain/clone have been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The deposited strains will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strains are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

30

Applicant's or agent's file reference KP/BM45332	International application
--	---------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>70</u> , line <u>3-28</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 UNIVERSITY BLVD, MANASSAS, VIRGINIA 20110-2209, UNITED STATES OF AMERICA	
Date of deposit 21 June 1997 (21.06.97) and 12 February 1999 (12.02.99)	Accession Number 43617 and 207099
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations where a European Patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn, only by issue of such a sample to an expert nominated by the person requesting the sample.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (If the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

BM45332

09/787083

CLAIMS:

JC08 Rec'd PCT/PTO 14 MAR 2001

1. An isolated polypeptide comprising an amino acid sequence which has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

3. The polypeptide as claimed in claim 1 comprising the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

4. An isolated polypeptide of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

5. An immunogenic fragment of the polypeptide as claimed in any one of claims 1 to 4 in which the immunogenic fragment is capable of raising an immune response (if necessary when coupled to carrier) which recognises the polypeptide of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

6. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least 85% identity to the amino acid sequence of SEQ ID NO:2, 4 6 or 8 over the entire length of SEQ ID NO:2, 4, 6 or 8 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

7. An isolated polynucleotide comprising a nucleotide sequence that has at least 85% identity to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2, 4, 6 or 8 over the

BM45322

entire coding region; or a nucleotide sequence complementary to said isolated polynucleotide.

8. An isolated polynucleotide which comprises a nucleotide sequence which has at least 85% identity to that of SEQ ID NO:1, 3, 5 or 7 over the entire length of SEQ ID NO:1, 3, 5 or 7 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

9. The isolated polynucleotide as claimed in any one of claims 6 to 8 in which the identity is at least 95% to SEQ ID NO:1, 3, 5 or 7.

10. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

11. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5 or SEQ ID NO:7.

12. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8, obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5 or SEQ ID NO:7 or a fragment thereof.

13. An expression vector comprising an isolated polynucleotide according to any one of claims 6-12.

14. A recombinant live microorganism comprising an expression vector according to claim 13.

15. A host cell comprising the expression vector of claim 13.

16. A recombinant membrane of the microorganism of claim 14 expressing an isolated polypeptide comprising an amino acid sequence that has at

AMENDED SHEET

8-09-2000

EP 009906781

BM45532

least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8.

17. A process for producing a polypeptide comprising an amino acid sequence that has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:8 comprising culturing a host cell of claim 15 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.
- 18.. A process for expressing a polynucleotide of any one of claims 6 - 12 comprising transforming a host cell with the expression vector comprising at least one of said polynucleotides and culturing said host cell under conditions sufficient for expression of any one of said polynucleotides.
19. A vaccine composition comprising an effective amount of the polypeptide of any one of claims 1 to 5 and a pharmaceutically acceptable carrier.
20. A vaccine composition comprising an effective amount of the polynucleotide of any one of claims 6 to 12 and a pharmaceutically effective carrier.
21. The vaccine composition according to either one of claims 19 or 20 wherein said composition comprises at least one other *Moraxella catarrhalis* antigen.
22. An antibody immunospecific for the polypeptide or immunological fragment as claimed in any one of claims 1 to 5.
23. A method of diagnosing a *Moraxella* infection, comprising identifying a polypeptide as claimed in any one of claims 1 - 5, or an antibody that is immunospecific for said

8-09-2000

BM45332

09/787083

JCO8 Rec'd PCT/PTO

14 MAR 2001

EP 009906781

polypeptide, present within a biological sample from an animal suspected of having such an infection.


24. Use of a composition comprising an immunologically effective amount of a polypeptide as claimed in any one of claims 1 - 5 in the preparation of a medicament for use in generating an immune response in an animal.

25. Use of a composition comprising an immunologically effective amount of a polynucleotide as claimed in any one of claims 6 - 12 in the preparation of a medicament for use in generating an immune response in an animal.

26. A therapeutic composition useful in treating humans with *Moraxella catarrhalis* disease comprising at least one antibody directed against the polypeptide of claims 1 - 5 and a suitable pharmaceutical carrier.

15


Figure 1 : Photographs of the electrophoresis gels obtained in the RFLP analysis of BASB034



WO 00/15802

PCT/EP99/06781

2/15



09727083 05440

WO 00/15802

3/15

PCT/EP99/06781





FIG. 10

WO 00/15802

PCT/EP99/06781

4/15



00787083 05440




Figure 2 : Alignment of the BASB034 polynucleotide sequences.

Identity to SeqID No:1 is indicated by a dot.

	*	20	*	40	*	
Seqid1 :	ATGAAAGTTTCACTGTCTACATTGACTTTATCTATTTTGT	CATGTTT	TGC :	50		
Seqid3 :	C.....	:	50		
Seqid5 :	:	50			
Seqid7 :	C.....	:	50		

	60	*	80	*	100	
Seqid1 :	TATCCTAGCCATT	CAGCAAGCACAAGCTGTACCAAATCCTGTGGCATT	TG :	100		
Seqid3 :	:	100			
Seqid5 :	A.....	:	100		
Seqid7 :	C.....	:	100			

	*	120	*	140	*	
Seqid1 :	TTGACGAAGTACGCAGTGAAAATGATCTTGGGCAAGACAATGAATTACCC	:	150			
Seqid3 :	A.....	T. :	150		
Seqid5 :	:	150			
Seqid7 :	:	150			

	160	*	180	*	200	
Seqid1 :	ATTGATGTCCAAAGT	GCGACACAATCAGCGTCTACTGATACGGCTAATCC	:	200		
Seqid3 :G...A.....	:	200			
Seqid5 :	:	200			
Seqid7 :	G.....	:	200		

	*	220	*	240	*	
Seqid1 :	TTTAGACGAACATGAACCAGAGCTTTATACGACAGCTTTAGAAAATAAAA	:	250			
Seqid3 :	:	250			
Seqid5 :	:	250			
Seqid7 :	:	250			

Publ. No. 00/15802

	*	320	*	340	*
Seqid1 :	TGCTATGACACTTTGGTGCATGGTGAGACGCCAGCGGTAATTTAAACCAA				: 350
Seqid3 :				: 350
Seqid5 :				: 350
Seqid7 :				: 350

```

          *          420          *          440          *
Seqid1 : CCCAGGTTATCTATCAAGAAACGACAGATCCGATTTTTTAAATGGGTAAT : 450
Seqid3 : .....G..... : 450
Seqid5 : .....G..... : 450
Seqid7 : .....G..... : 450

```

Seqid1 : CAAACAGTTTACACCACTGAGCTTATCATTTGATTTAGACCGAAATAATA : 550

8/15

Seqid3 : : 550
Seqid5 : : 550
Seqid7 : : 550

560 * 580 * 600
Seqid1 : CACCACTTTGGTCATCAGACCACACAATCCGATGTATGTATTGCCCATA : 600
Seqid3 :G..... : 600
Seqid5 : : 600
Seqid7 : : 600

* 620 * 640 *
Seqid1 : TTTATGCACGGTAAGCCTAATCGAAGCCCAAATACGCCCAGTCATGAAGC : 650
Seqid3 : : 650
Seqid5 : : 650
Seqid7 : : 650

660 * 680 * 700
Seqid1 : AAAACAATTTACCCCAAATGAATTTTCGTGCTCCCGAGCTAAAATTTTCAGG : 700
Seqid3 : ..G.....C..T..AT.....A. : 700
Seqid5 : ..G.....C..T..AT.....A. : 700
Seqid7 : : 700

* 720 * 740 *
Seqid1 : TTTCTGTTAAGGTTAAAGCTGCTGAGGATTATGGGGGACGGATTCAGAT : 750
Seqid3 : : 750
Seqid5 : : 750
Seqid7 : : 750

760 * 780 * 800
Seqid1 : TTATGGTTTGGATATACACAGCAATCGCACTGGCAGATTTTAAATGGAAA : 800
Seqid3 :G..... : 800
Seqid5 : : 800
Seqid7 : : 800

WO 00/15802

PCT/EP99/06781

9/15

* 820 * 840 *

Seqid1 : AAACCTCTCGTCCTTTTAGAGTACATGACTACCAGCCAGAGATTTTCTTAA : 850

Seqid3 :T..... : 850

Seqid5 :T..... : 850

Seqid7 : : 850

 860 * 880 * 900

Seqid1 : CTCAACCTGTATACTCAGACTTACCATGGGATGGCAAAGTCCGCATGATT : 900

Seqid3 :G..... : 900

Seqid5 : : 900

Seqid7 : : 900

 * 920 * 940 *

Seqid1 : GGCATGGGTGCGGTACATCATTCGAATGGTGAAAGTGCCAAACTGTCTCG : 950

Seqid3 : : 950

Seqid5 : : 950

Seqid7 : : 950

 960 * 980 * 1000

Seqid1 : CTCATGGAATCGTGCTTATTTGATGGCAGGCATGGAATGGAAAAACCTGA : 1000

Seqid3 : : 1000

Seqid5 : : 1000

Seqid7 : : 1000

 * 1020 * 1040 *

Seqid1 : CTGTCATGCCACGCATTTGGGGCGTATCTTTAAAGAGGGTAGTGGCAGC : 1050

Seqid3 : : 1050

Seqid5 : : 1050

Seqid7 : : 1050

 1060 * 1080 * 1100

Seqid1 : CAGCCAGATGATAATCCTGATATCTTGACTATTATGGTTATGGTGATGT : 1100

Seqid3 :C..... : 1100

Seqid5 :C..... : 1100

09/787083 051444

10/15

Seqid7 : : 1100

* 1120 * 1140 *

Seqid1 : GCGTTTTTTATATCAACTAGAAAATAAAAGTAATATTTTCAGGTACGGTAC : 1150

Seqid3 : : 1150

Seqid5 : : 1150

Seqid7 : : 1150

1160 * 1180 * 1200

Seqid1 : GCTATAATCCACGCTCAGGCCAAAGGTGCGTTGCAACTTGACTATGTCTAT : 1200

Seqid3 : : 1200

Seqid5 : : 1200

Seqid7 : : 1200

* 1220 * 1240 *

Seqid1 : CCGCTTGGTAAGGGAATTAGTGGCTATTTTCAAATATTTCAAGGCTATGG : 1250

Seqid3 : : 1250

Seqid5 : : 1250

Seqid7 : : 1250

1260 * 1280 * 1300

Seqid1 : GCAGTCTTTGATTGATTATAATCATGAGGCGACAAGCTTTGGCGTCGGAC : 1300

Seqid3 : : 1300

Seqid5 : : 1300

Seqid7 : : 1300

* 1320

Seqid1 : TTATGCTTAACGACTGGATGGGTCTATAA : 1329

Seqid3 : : 1329

Seqid5 : : 1329

Seqid7 : : 1329

T 004753 004753 004753

11/15

Figure 3 : Alignment of the BASB034 polypeptide sequences.

Identity to SeqID No:2 is indicated by a dot.

```

          *      20      *      40      *
Seqid2 : MKVSLSTLTLSILSCFAILAIQQAQAVPNPVAFAVDEVRSNDLGQDNELP : 50
Seqid4 : .....P.....K.....L : 50
Seqid6 : .....K..... : 50
Seqid8 : .....P..... : 50

          60      *      80      *      100
Seqid2 : IDVQSATQSASTDTANPLDEHEPELYTTALENKTMLINCSALNQDIMRLA : 100
Seqid4 : .G..... : 100
Seqid6 : ..... : 100
Seqid8 : ..... : 100

          *      120      *      140      *
Seqid2 : CYDTLVHGETPAVIKTKRSIRLDETIWQTIKGKPQVIYQETTDPIFLMGN : 150
Seqid4 : .....V..... : 150
Seqid6 : .....V..... : 150
Seqid8 : .....V..... : 150

          160      *      180      *      200
Seqid2 : EKGMLTKKDAKQLEYAAKQFTPLSLSFDLDRNNTPLWSSRPHNPMYVLPI : 200
Seqid4 : ..... : 200
Seqid6 : ..... : 200
Seqid8 : ..... : 200

          *      220      *      240      *
Seqid2 : FMHGKPNRSPNTPSHEAKQFTPNEFRAPELKFQVSVKVKAAEDLWGTDS : 250
Seqid4 : .....R..... : 250
Seqid6 : .....R..... : 250
Seqid8 : ..... : 250

```

260 * 280 * 300
Seqid2 : LWFGYTQQSHWQIFNGKNSRPFVRVHDYQPEIFLTQPVYSDLPWDGKVRMI : 300
Seqid4 : : 300
Seqid6 : : 300
Seqid8 : : 300


 * 320 * 340 *
Seqid2 : GMGAVHHSNGESAKLSRSWNRAYLMAGMEWKNLTVMPRIWGRIFKEGSGS : 350
Seqid4 : : 350
Seqid6 : : 350
Seqid8 : : 350

 360 * 380 * 400
Seqid2 : QPDDNPDILDYYGYGDVRFYQLENKSNISGTVRYNPRSGKALQLDYVY : 400
Seqid4 : : 400
Seqid6 : : 400
Seqid8 : : 400

 * 420 * 440
Seqid2 : PLGKGISGYFQIFQGYGQSLIDYNHEATSFGVGLMLNDWMGL : 442
Seqid4 : : 442
Seqid6 : : 442
Seqid8 : : 442

Publ. No. WO 00/15802

Figure 4 : Coomassie stained SDS-PAGE of recombinant purified BASB034



14/15

Figure 5 : Western-blot of purified recombinant BASB034 protein probed with anti-His antibody





Figure 6 : Western-blot of BASB034 probed with immune (membrane 1; right lane) and pre-immune (membrane 2; right lane) rabbit sera at 1:500. The arrow indicates the position of BASB034 on the membrane. The left lanes on membrane 1 and 2 are molecular weight markers.

[illegible]

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

MORAXELLA CATARRHALIS BASB034 POLYPEPTIDES AND USES THEREOF

the specification of which (check one)

☐ is attached hereto.

☒ was filed on 14 September 1999 as Serial No. PCT/EP99/06781
and was amended on (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or Inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Number	Country	Filing Date	Priority Claimed
9820002.5	Great Britain	14 September 1998	Yes

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

Application Number	Filing Date
--------------------	-------------

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

I hereby appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

Customer Number 25,308

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of Inventor: Jean-Louis RUELLE

Inventor's Signature: _____

Date: March 21st, 2002

Residence: LIMAL, BELGIUM

Citizenship: BELGIAN

Post Office Address: SmithKline Beecham Corporation
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939

SEQUENCE LISTING

<110> SmithKline Beecham Biologicals S.A.

<120> Novel Compounds

<130> BM45332

<160> 13

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1329

<212> DNA

<213> *Moraxella catarrhalis*

<400> 1

atgaaagttt cactgtctac attgacttta tctattttgt catgttttgc taccctagcc	60
attcagcaag cacaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtga	120
aatgatcttg ggcaagacaa tgaattaccc attgatgtcc aaagtgcgac acaatcagcg	180
tctactgata cggctaattcc tttagacgaa catgaaccag agctttatac gacagcttta	240
gaaaataaaa ccatgctgat taactgtcca gcaacttaac aagatatcat gcgtttggcg	300
tgctatgaca ctttgggtgca tgggtgagacg ccagcggtaa ttaaaaccaa gcgttccatt	360
cgccttgatg aaacaatttg gcagaccatc aaaggcaaac cccaggttat ctatcaagaa	420
acgacagatc cgattttttt aatgggtaac gaaaaaggca tgctgaccaa aaaagatgcc	480
aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac	540
cgaaataata caccactttg gtcattcaga ccacacaatc cgatgtatgt attgcccata	600
tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcattgaagc aaaacaattt	660
accccaaatg aatttcgtgc tcccagccta aaatttcagg tttctgttaa gggttaaagct	720
gctgaggatt tatgggggac ggattcagat ttatgggttg gatatacaca gcaatcgcac	780
tggcagattt ttaatggaaa aaactctcgt ccttttagag tacatgacta ccagccagag	840
attttcttaa ctcaacctgt atactcagac ttaccatggg atggcaaatg ccgcatgatt	900
ggcatgggtg cggtagatca tcccaatggg gaaagtgcc aactgtctcg ctcatggaat	960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg	1020
gggcgtatct ttaagagggt tagtggcagc cagccagatg ataactctga tatcttggac	1080
tactatgggt atggctgatg gcgtttttta tatcaactag aaaataaaag taatattcca	1140

ggtagcggtac gctataatcc acgctcaggc aaaggtgcgt tgcaacttga ctatgtctat 1200
 ccgcttggtta aggggaattag tggctatttt caaatatttc aaggctatgg gcagtctttg 1260
 attgattata atcatgaggc gacaagcttc ggcgtcggac ttatgcttaa cgactggatg 1320
 ggtctataa 1329

<210> 2

<211> 442

<212> PRT

<213> *Moraxella catarrhalis*

<400> 2

Met	Lys	Val	Ser	Leu	Ser	Thr	Leu	Thr	Leu	Ser	Ile	Leu	Ser	Cys	Phe
1				5					10					15	
Ala	Ile	Leu	Ala	Ile	Gln	Gln	Ala	Gln	Ala	Val	Pro	Asn	Pro	Val	Ala
				20				25						30	
Phe	Val	Asp	Glu	Val	Arg	Ser	Glu	Asn	Asp	Leu	Gly	Gln	Asp	Asn	Glu
				35				40						45	
Leu	Pro	Ile	Asp	Val	Gln	Ser	Ala	Thr	Gln	Ser	Ala	Ser	Thr	Asp	Thr
				50				55						60	
Ala	Asn	Pro	Leu	Asp	Glu	His	Glu	Pro	Glu	Leu	Tyr	Thr	Thr	Ala	Leu
65						70				75					80
Glu	Asn	Lys	Thr	Met	Leu	Ile	Asn	Cys	Ser	Ala	Leu	Asn	Gln	Asp	Ile
				85						90					95
Met	Arg	Leu	Ala	Cys	Tyr	Asp	Thr	Leu	Val	His	Gly	Glu	Thr	Pro	Ala
				100						105					110
Val	Ile	Lys	Thr	Lys	Arg	Ser	Ile	Arg	Leu	Asp	Glu	Thr	Ile	Trp	Gln
				115						120					125
Thr	Ile	Lys	Gly	Lys	Pro	Gln	Val	Ile	Tyr	Gln	Glu	Thr	Thr	Asp	Pro
				130						135					140
Ile	Phe	Leu	Met	Gly	Asn	Glu	Lys	Gly	Met	Leu	Thr	Lys	Lys	Asp	Ala
145						150				155					160
Lys	Gln	Leu	Glu	Tyr	Ala	Ala	Lys	Gln	Phe	Thr	Pro	Leu	Ser	Leu	Ser
				165						170					175
Phe	Asp	Leu	Asp	Arg	Asn	Asn	Thr	Pro	Leu	Trp	Ser	Ser	Arg	Pro	His
				180						185					190
Asn	Pro	Met	Tyr	Val	Leu	Pro	Ile	Phe	Met	His	Gly	Lys	Pro	Asn	Arg
				195						200					205
Ser	Pro	Asn	Thr	Pro	Ser	His	Glu	Ala	Lys	Gln	Phe	Thr	Pro	Asn	Glu
				210						215					220

WO 00/15802

PCT/EP99/06781

Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
 225 230 235 240
 Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
 245 250 255
 Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
 260 265 270
 Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
 275 280 285
 Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala
 290 295 300
 Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
 305 310 315 320
 Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
 325 330 335
 Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
 340 345 350
 Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
 355 360 365
 Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
 370 375 380
 Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
 385 390 395 400
 Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
 405 410 415
 Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
 420 425 430
 Gly Leu Met Leu Asn Asp Trp Met Gly Leu
 435 440

<210> 3

<211> 1329

<212> DNA

<213> *Moraxella catarrhalis*

<400> 3

atgaaagttt cactgtctac attgacttta tctattttgc catgttttgc tctcctagcc	60
attcagcaag cacaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtaaa	120
aatgatcttg ggcaagacaa tgaattactc attggtgtac aaagtgcgac acaatcagcg	180
tctactgata cggctaatacc tttagacgaa catgaaccag agctttatag gacagcttta	240

gaaaataaaaa ccatgctgat taactgctca gcacttaatc aagatatcat gcgtttggcg 300
 tgctatgaca ctttgggtgca tggtgagacg ccagcggtaa ttaaaaccaa gcgttccatt 360
 cgccttgatg aaacaatttg gcagaccatc aaaggcaaac cccaggttgt ctatcaagaa 420
 acgacagatc cgattttttt aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc 480
 aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatc tgatttagac 540
 cgaaataata caccgctttg gtcacacga ccacacaatc cgatgtatgt attgcccata 600
 tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcacgaagc aagacaattt 660
 accccaaatg aatttcgtgc ccctgaatta aaatttcaag tttctgttaa gggttaaagct 720
 gctgaggatt tatgggggac ggattcagat ttatgggttg ggtatacaca gcaatcgac 780
 tggcagattt ttaatggaaa aaactctcgt ccttttagag tacatgatta ccagccagag 840
 attttcttaa ctcaacctgt gtactcagac ttaccatggg atggcaaagt cgcgatgatt 900
 ggcacgggtg cggtacatca ttccaatggt gaaagtgcc aactgtctcg ctcatggaat 960
 cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg 1020
 gggcgatctt ttaaagaggg tagtggcagc cagccagatg acaatcctga tatcttgac 1080
 tattatgggt atgggtgatg gcgtttttta tatcaactag aaaataaaag taatatttca 1140
 ggtacgggtac gctataatcc acgctcaggc aaaggtgcgt tgcaacttga ctatgtctat 1200
 ccgcttggtg aggggaattag tggctatttt caaatatttc aaggctatgg gcagcttttg 1260
 attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg 1320
 ggtctataa 1329

<210> 4

<211> 442

<212> PRT

<213> *Moraxella catarrhalis*

<400> 4

Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Pro Cys Phe
 1 5 10 15
 Ala Ile Leu Ala Ile Gln Gln Ala Gln Ala Val Pro Asn Pro Val Ala
 20 25 30
 Phe Val Asp Glu Val Arg Ser Lys Asn Asp Leu Gly Gln Asp Asn Glu
 35 40 45
 Leu Leu Ile Gly Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
 50 55 60
 Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
 65 70 75 80
 Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
 85 90 95
 Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala

100 105 110
 Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
 115 120 125
 Thr Ile Lys Gly Lys Pro Gln Val Val Tyr Gln Glu Thr Thr Asp Pro
 130 135 140
 Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
 145 150 155 160
 Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
 165 170 175
 Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
 180 185 190
 Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
 195 200 205
 Ser Pro Asn Thr Pro Ser His Glu Ala Arg Gln Phe Thr Pro Asn Glu
 210 215 220
 Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
 225 230 235 240
 Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
 245 250 255
 Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
 260 265 270
 Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
 275 280 285
 Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala
 290 295 300
 Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
 305 310 315 320
 Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
 325 330 335
 Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
 340 345 350
 Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
 355 360 365
 Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
 370 375 380
 Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
 385 390 395 400
 Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
 405 410 415

Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val

420

425

430

Gly Leu Met Leu Asn Asp Trp Met Gly Leu

435

440

<210> 5

<211> 1329

<212> DNA

<213> *Moraxella catarrhalis*

<400> 5

```

atgaaagttt cactgtctac attgacttta tctattttgt catgttttgc taccctagcc      60
attcagcaag caaaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtgaa      120
aatgatcttg ggcaagacaa tgaattaccc attgatgtcc aaagtgcgac acaatcagcg      180
tctactgata cggctaatacc tttagacgaa catgaaccag agctttatac gacagcttta      240
gaaaataaaa ccatgctgat taactgctca gcacttaatc aagatatcat gcgtttggcg      300
tgctatgaca ctttgggtgca tgggtgagac ccagcggtaa ttaaaaccaa gcgttccatt      360
cgccttgatg aaacaatttg gcagaccatc aaaggcaaac cccaggttgt ctatcaagaa      420
acgacagatc cgattttttt aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc      480
aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac      540
cgaaataata caccactttg gtcatcacga ccacacaatc cgatgtatgt attgcccata      600
tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcatgaagc aagacaattt      660
accccaaatg aatttcgtgc ccctgaatta aaatttcaag tttctgttaa gggttaaagct      720
gctgaggatt tatgggggac ggattcagat ttatggtttg gatatacaca gcaatcgcac      780
tggcagattt ttaatggaaa aaactctcgt ctttttagag tacatgatta ccagccagag      840
attttcttaa ctcaacctgt atactcagac ttaccatggg atggcaaatg ccgcatgatt      900
ggcatgggtg cggtagatca ttccaatggc gaaagtgcc aactgtctcg ctcatggaat      960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg      1020
gggcgtatct ttaaagaggg tagtggcagc cagccagatg acaatcctga tatcttggac      1080
tattatggtt atggtgatgt gcgtttttta tatcaactag aaaataaaag taatatttca      1140
ggtacggtag gctataatcc acgctcaggc aaagggtcgt tgcaacttga ctatgtctat      1200
ccgcttggta agggaattag tggctatttt caaatatttc aaggctatgg gcagtccttg      1260
attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg      1320
ggtctataa                                     1329

```

<210> 6

<211> 442

<212> PRT

<213> *Moraxella catarrhalis*

<400> 6

```

Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Ser Cys Phe
  1             5             10             15
Ala Ile Leu Ala Ile Gln Gln Ala Lys Ala Val Pro Asn Pro Val Ala
      20             25             30
Phe Val Asp Glu Val Arg Ser Glu Asn Asp Leu Gly Gln Asp Asn Glu
      35             40             45
Leu Pro Ile Asp Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
      50             55             60
Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
      65             70             75             80
Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
      85             90             95
Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala
      100            105            110
Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
      115            120            125
Thr Ile Lys Gly Lys Pro Gln Val Val Tyr Gln Glu Thr Thr Asp Pro
      130            135            140
Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
      145            150            155            160
Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
      165            170            175
Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
      180            185            190
Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
      195            200            205
Ser Pro Asn Thr Pro Ser His Glu Ala Arg Gln Phe Thr Pro Asn Glu
      210            215            220
Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
      225            230            235            240
Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
      245            250            255
Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
      260            265            270
Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
      275            280            285
Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala

```

290 295 300
 Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
 305 310 315 320
 Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
 325 330 335
 Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
 340 345 350
 Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
 355 360 365
 Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
 370 375 380
 Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
 385 390 395 400
 Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
 405 410 415
 Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
 420 425 430
 Gly Leu Met Leu Asn Asp Trp Met Gly Leu
 435 440

<210> 7

<211> 1329

<212> DNA

<213> *Moraxella catarrhalis*

<400> 7

atgaaagttt cactgtctac attgacttta tctatttttg catgttttgc catcctagcc 60
 attcagcaag cacaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtga 120
 aatgatcttg ggcaagacaa tgaattaccc attgatgtcc aaagtgcgac acaatcggcg 180
 tctactgata cggctaattc tttagacgaa catgaaccag agctttatac gacagcttta 240
 gaaaataaaa ccatgctgat taactgtcca gcacttaatc aagatatcat gcgtttggcg 300
 tgctatgaca ctttggtgca tgggtgagac ccagcggtaa ttaaaaccaa gcgtttccatt 360
 cgccttgatg aaacaatttg gcagaccatc aaaggcaaac cccagggttg ctatcaagaa 420
 acgacagatc cgattttttt aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc 480
 aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac 540
 cgaaataata caccactttg gtcattcaga ccacacaatc cgatgtatgt attgcccata 600
 tttatgcacg gtaagcctaa tcgaagccca aatcgcacca gtcattgaagc aaaacaattt 660
 accccaaatg aatttcgtgc tcccagacta aaatttcagg tttctgttaa gggttaaagct 720
 gctgaggatt tatggggggac ggattcagat ttatgggttg gatatacaca gcaatcgcac 780

```

tggcagattt ttaatggaaa aaactctcgt ccttttagag tacatgacta ccagccagag      840
attttcttaa ctcaacctgt atactcagac ttaccatggg atggcaaagt ccgcatgatt      900
ggcatgggtg cggtagatca ttccaatggt gaaagtgcc aactgtctcg ctcattggaat     960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg    1020
gggcgtatct ttaaagaggg tagtggcagc cagccagatg ataatcctga tatcttggac    1080
tattatgggt atggtgatgt gcgtttttta tatcaactag aaaataaaaag taatatttca    1140
ggcacgggtac gctataatcc acgctcaggc aaaggtgcgt tgcaacttga ctatgtctat    1200
ccgcttggtg agggaattag tggctatttt caaatatttc aaggctatgg gcagtctttg    1260
attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg    1320
ggctctataa                                     1329

```

<210> 8

<211> 442

<212> PRT

<213> *Moraxella catarrhalis*

<400> 8

```

Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Pro Cys Phe
 1             5             10            15
Ala Ile Leu Ala Ile Gln Gln Ala Gln Ala Val Pro Asn Pro Val Ala
      20             25            30
Phe Val Asp Glu Val Arg Ser Glu Asn Asp Leu Gly Gln Asp Asn Glu
      35             40            45
Leu Pro Ile Asp Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
      50             55            60
Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
      65             70            75            80
Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
      85             90            95
Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala
      100            105            110
Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
      115            120            125
Thr Ile Lys Gly Lys Pro Gln Val Val Tyr Gln Glu Thr Thr Asp Pro
      130            135            140
Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
      145            150            155            160
Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
      165            170            175

```


Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
 180 185 190
 Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
 195 200 205
 Ser Pro Asn Thr Pro Ser His Glu Ala Lys Gln Phe Thr Pro Asn Glu
 210 215 220
 Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
 225 230 235 240
 Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
 245 250 255
 Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
 260 265 270
 Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
 275 280 285
 Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala
 290 295 300
 Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
 305 310 315 320
 Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
 325 330 335
 Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
 340 345 350
 Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
 355 360 365
 Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
 370 375 380
 Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
 385 390 395 400
 Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
 405 410 415
 Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
 420 425 430
 Gly Leu Met Leu Asn Asp Trp Met Gly Leu
 435 440

<210> 9

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer sequence

<400> 9

gatttaagag tatgttatga tg

22

<210> 10

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer sequence

<400> 10

gtatgggttg atcaaataca g

21

<210> 11

<211> 58

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide

<400> 11

aagggcccaa ttacgcagag gggatcccaa gctgtaccaa atcctgtggc atttgttg

58

<210> 12

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide

<400> 12

aagggcccaa ttacgcagag ggtcgactta ttatagacc atccagtcgt taagcataag

60

<210> 13

<211> 1000

<212> DNA

<213> *Moraxella catarrhalis*

<400> 13

```

acttgccgaa aataccatctt atactgattg tgatgttata caggcagatg gcggtacacg      60
cacagccagt atcagtgggtg ctgcggtggc acttattgat gctttagaac acttgacagc      120
tcgtaaaaag cttacccaag atccgctttt gggcttggtg gcagcgggtt ctgtgggtgt      180
taatcaaggc cgtgtattgc ttgatttggg ttatgctgaa gattcaactt gtgataccga      240
tttaaatgtg gtcattgacgc aggcaggtgg gtttattgag attcaaggca cagcagaaga      300
aaagccatctt actcgtgctg aagctaattg gatgcttgat ttggcagagc tgggaattgg      360
gcagattatc gaagcccaaa agcaagtatt aggcctggta tatgctaata gttgaagata      420
atggcgtgat catcacatta aatggacaag taaaagaccc attatttttg tggtcgatga      480
tattgctgct gctgggtgtc ttgggtggca tcatttgttt gattgcaccc gttttttatg      540
caatcgggtg gttggcctta ttgcagttg tggattttgt gtttaatat caaaggcaaa      600
aagccaaaac ttgtcatatg ttttcacaag gtcgcttgaa gattacgtcc aaacgctttg      660
agattcataa caaatcacta accttatcag catcggcaac aatatctgct aaagataaca      720
aaatgacaat tgttgatcgg ggcattgaat atcattttac aggttttgct gatgaccgtg      780
aaattaatat agccaaacag gtacttttgg gaaagtcaat caaaaccaat gcggtggcgg      840
taacattggc taagtagttg ttgtgataca gacaggttgg atggctctta actccacca      900
cctaactttt tctttgtttg gatttaagag tatgttatga tgggcaggat tttattttaa      960
gtcatcattt aatgcaatca gttgtccaga gtagccgttc      1000

```

09/787083

JGUB Rec'd PCT/PTO 14 MAR 2001

SEQUENCE LISTING

<110> Ruelle, Jean-Louis

<120> Novel Compounds

<130> BM45332

<150> PCT/EP99/06781

<151> 1999-09-14

<150> GB 9820002.5

<151> 1998-09-14

<160> 13

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1329

<212> DNA

<213> Moraxella catarrhalis

<400> 1

```

atgaaagttt cactgtctac attgacttta tctattttgt catgttttgc tctcctagcc      60
attcagcaag cacaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtga      120
aatgatcttg ggcaagacaa tgaattaccc attgatgtcc aaagtgcgac acaatcagcg      180
tctactgata cggctaatacc tttagacgaa catgaaccag agctttatac gacagcttta      240
gaaaataaaa ccattgctgat taactgctca gcacttaatc aagatatcat gcgtttggcg      300
tgctatgaca ctttgggtgca tggtgagacg ccagcggtaa ttaaaaccaa gcgttccatt      360
cgctttgatg aaacaatttg gcagaccatc aaaggcaaac cccaggttat ctatcaagaa      420
acgacagatc cgattttttt aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc      480
aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac      540
cgaaataata caccactttg gtcatcacga ccacacaatc cgatgtatgt attgcccata      600
tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcatgaagc aaaacaattt      660
accccaaatg aatttcgtgc tcccgagcta aaatttcagg tttctgttaa ggttaaagct      720
gctgaggatt tatgggggac ggattcagat ttatggtttg gatatacaca gcaatcgac      780
tggcagattt ttaatggaaa aaactctcgt ccttttagag tacatgacta ccagccagag      840
attttcttaa ctcaacctgt atactcagac ttaccatggg atggcaaagt ccgcatgatt      900
ggcatgggtg cggtacatca ttccaatggt gaaagtgcc aactgtctcg ctcatggaat      960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg     1020
gggcgtatct ttaaagaggg tagtggcagc cagccagatg ataactctga tatcttggac     1080
tattatggtt atgggtgatg gcgtttttta tatcaactag aaaataaaaag taatatttca     1140
ggtacggtac gctataatcc acgctcaggc aaagggtgcgt tgcaacttga ctatgtctat     1200
ccgcttggtg aggggaattag tggctatttt caaatatttc aaggctatgg gcagtctttg     1260
attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg     1320
ggtctataa                                     1329

```

<210> 2

<211> 442

<212> PRT

<213> Moraxella catarrhalis

<400> 2

```

Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Ser Cys Phe
 1             5             10             15
Ala Ile Leu Ala Ile Gln Gln Ala Gln Ala Val Pro Asn Pro Val Ala
                20                25                30
Phe Val Asp Glu Val Arg Ser Glu Asn Asp Leu Gly Gln Asp Asn Glu
          35                   40                   45

```

```

Leu Pro Ile Asp Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
 50                               55                               60
Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
 65                               70                               75                               80
Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
                               85                               90                               95
Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala
                               100                            105                            110
Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
                               115                            120                            125
Thr Ile Lys Gly Lys Pro Gln Val Ile Tyr Gln Glu Thr Thr Asp Pro
                               130                            135                            140
Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
145                               150                               155                               160
Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
                               165                               170                               175
Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
                               180                            185                            190
Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
                               195                            200                            205
Ser Pro Asn Thr Pro Ser His Glu Ala Lys Gln Phe Thr Pro Asn Glu
                               210                            215                            220
Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
225                               230                               235                               240
Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
                               245                               250                               255
Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
                               260                            265                            270
Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
                               275                            280                            285
Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala
290                               295                               300
Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
305                               310                               315                               320
Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
                               325                            330                            335
Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
                               340                            345                            350
Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
                               355                            360                            365
Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
370                               375                               380
Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
385                               390                               395                               400
Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
                               405                               410                               415
Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
                               420                            425                            430
Gly Leu Met Leu Asn Asp Trp Met Gly Leu
                               435                               440

```

<210> 3

<211> 1329

<212> DNA

<213> Moraxella catarrhalis

<400> 3

```

atgaaagttt cactgtctac attgacttta tctatatttgc catgtttttgc tctcctagcc      60
attcagcaag cacaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtaaa      120
aatgatcttg ggcaagacaa tgaattactc attgggtgtac aaagtgcgac acaatcagcg      180
tctactgata cggctaatacc tttagacgaa catgaaccag agctttatac gacagcttta      240
gaaaataaaa ccatgctgat taactgctca gcacttaatc aagatatcat gcggttggcg      300

```

```

tgctatgaca ctttggtgca tgggtgagacg ccagcggttaa ttaaaaccaa gcgttccatt 360
cgcccttgatg aaacaatttg gcagaccatc aaaggcaaac cccaggttgt ctatcaagaa 420
acgacagatc cgatTTTTTT aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc 480
aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac 540
cgaaataata caccgctttg gtcacacga ccacacaatc cgatgtatgt attgcccata 600
tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcatagaagc aagacaattt 660
accccaaatg aatttcgtgc ccctgaatta aaatttcaag tttctgttaa ggtaaagct 720
gctgaggatt tatgggggac ggattcagat ttatggtttg ggtatacaca gcaatcgcac 780
tggcagatTTT ttaatggaaa aaactctcgt ccttttagag tacatgatta ccagccagag 840
atTTTcttaa ctcaacctgt gtactcagac ttaccatggg atggcaaagt ccgcatgatt 900
ggcatgggtg cggtacatca ttccaatggg gaaagtgcc aactgtctcg ctcatggaat 960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg 1020
gggCGtatct ttaaagaggg tagtggcagc cagccagatg acaatcctga tatcttgac 1080
tattatggtt atgggtgatgt gcgtTTTTTT tatcaactag aaaataaaaag taatatttca 1140
ggtacgggtac gctataatcc acgctcaggc aaagggtcgt tgcaacttga ctatgtctat 1200
cgccttggtg aggggaattag tggctatTTT caaatatttc aaggctatgg gcagtctttg 1260
attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg 1320
ggtctataa 1329

```

<210> 4

<211> 442

<212> PRT

<213> Moraxella catarrhalis

<400> 4

```

Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Pro Cys Phe
  1          5          10          15
Ala Ile Leu Ala Ile Gln Gln Ala Gln Ala Val Pro Asn Pro Val Ala
  20          25          30
Phe Val Asp Glu Val Arg Ser Lys Asn Asp Leu Gly Gln Asp Asn Glu
  35          40          45
Leu Leu Ile Gly Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
  50          55          60
Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
  65          70          75          80
Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
  85          90          95
Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala
 100          105          110
Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
 115          120          125
Thr Ile Lys Gly Lys Pro Gln Val Val Tyr Gln Glu Thr Thr Asp Pro
 130          135          140
Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
 145          150          155          160
Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
 165          170          175
Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
 180          185          190
Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
 195          200          205
Ser Pro Asn Thr Pro Ser His Glu Ala Arg Gln Phe Thr Pro Asn Glu
 210          215          220
Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
 225          230          235          240
Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
 245          250          255
Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
 260          265          270
Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
 275          280          285
Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala

```

```

      290      295      300
Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
305      310      315      320
Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
      325      330      335
Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
      340      345      350
Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
      355      360      365
Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
      370      375      380
Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
385      390      395      400
Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
      405      410      415
Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
      420      425      430
Gly Leu Met Leu Asn Asp Trp Met Gly Leu
      435      440

```

```

<210> 5
<211> 1329
<212> DNA
<213> Moraxella catarrhalis

```

```

<400> 5
atgaaagttt cactgtctac attgacttta tctattttgt catgttttgc tctcctagcc      60
attcagcaag caaaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagtgaa      120
aatgatcttg ggcaagacaa tgaattacc attgatgtcc aaagtgcgac acaatcagcg      180
tctactgata cggctaatac ttttagacgaa catgaaccag agctttatac gacagcttta      240
gaaaataaaa ccatgctgat taactgctca gcacttaatc aagatatcat gcgtttggcg      300
tgctatgaca ctttggtgca tgggtgagacg ccagcggtaa ttaaaaccaa gcgttccatt      360
cgcccttgatg aaacaatttg gcagaccatc aaaggcgaac cccaggttgt ctatcaagaa      420
acgacagatc cgattttttt aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc      480
aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac      540
cgaaataata caccactttg gtcatacaga ccacacaatc cgatgtatgt attgcccata      600
tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcatagaagc aagacaattt      660
accccaaagt aatttcgtgc ccctgaatta aaatttcaag tttctgttaa gggttaaagct      720
tgctgaggatt tatgggggac ggattcagat ttatggtttg gatatacaca gcaatcgcac      780
tggcagattt ttaatggaat aaactctcgt ccttttagag tacatgatta ccagccagag      840
attttcttaa ctcaacctgt atactcagac ttaccatggg atggcaaagt ccgcatgatt      900
ggcatgggtg cggtaacatc ttccaatggg gaaagtgcc aactgtctcg ctcatggaat      960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg      1020
gggcgtatct ttaaagaggg tagtggcagc cagccagatg acaatcctga tatcttggac      1080
tattatgggtt atgggtgatg gcgtttttta tatcaactag aaaataaaaag taatatttca      1140
ggtacgggtac gctataatcc acgctcaggc aaagggtcgt tgcaacttga ctatgtctat      1200
ccgcttggtg agggaattag tggctatitt caaatatttc aaggctatgg gcagtctttg      1260
attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg      1320
ggtctataa
      1329

```

```

<210> 6
<211> 442
<212> PRT
<213> Moraxella catarrhalis

```

```

<400> 6
Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Ser Cys Phe
1      5      10      15
Ala Ile Leu Ala Ile Gln Gln Ala Lys Ala Val Pro Asn Pro Val Ala
20      25      30
Phe Val Asp Glu Val Arg Ser Glu Asn Asp Leu Gly Gln Asp Asn Glu
35      40      45

```

```

Leu Pro Ile Asp Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
  50                      55                      60
Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
 65                      70                      75                      80
Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
                      85                      90                      95
Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala
                      100                    105                    110
Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
                      115                    120                    125
Thr Ile Lys Gly Lys Pro Gln Val Val Tyr Gln Glu Thr Thr Asp Pro
                      130                    135                    140
Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
145                      150                      155                      160
Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
                      165                      170                      175
Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
                      180                    185                    190
Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
                      195                    200                    205
Ser Pro Asn Thr Pro Ser His Glu Ala Arg Gln Phe Thr Pro Asn Glu
                      210                    215                    220
Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
225                      230                      235                      240
Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
                      245                    250                    255
Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
                      260                    265                    270
Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
                      275                    280                    285
Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala
290                      295                      300
Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
305                      310                      315                      320
Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
                      325                    330                    335
Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
                      340                    345                    350
Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
                      355                    360                    365
Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
                      370                    375                    380
Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
385                      390                      395                      400
Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
                      405                    410                    415
Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
                      420                    425                    430
Gly Leu Met Leu Asn Asp Trp Met Gly Leu
                      435                    440

```

<210> 7

<211> 1329

<212> DNA

<213> Moraxella catarrhalis

<400> 7

```

atgaaagttt cactgtctac attgacttta tctattttgc catgttttgc catcctagcc      60
attcagcaag cacaagctgt accaaatcct gtggcatttg ttgacgaagt acgcagttaa      120
aatgatcttg ggcaagacaa tgaattaccc attgatgtcc aaagtgcgac acaatcggcg      180
tctactgata cggctaattc tttagacgaa catgaaccag agctttatac gacagcttta      240
gaaaataaaa ccatgctgat taactgctca gcacttaatc aagatatcat gcgtttggcg      300

```



```

tgctatgaca ctttggtgca tgggtgagacg ccagcggtaa ttaaaaccaa gcgttccatt 360
cgccttgatg aaacaatttg gcagaccatc aaaggcaaac cccaggttgt ctatcaagaa 420
acgacagatc cgattttttt aatgggtaat gaaaaaggca tgctgaccaa aaaagatgcc 480
aaacagcttg aatatgcagc caaacagttt acaccactga gcttatcatt tgatttagac 540
cgaaataata caccactttg gtcatcacga ccacacaatc cgatgtatgt attgcccata 600
tttatgcacg gtaagcctaa tcgaagccca aatacgccca gtcatgaagc aaaacaattt 660
accccaaatg aatttcgtgc tcccgagcta aaatttcagg tttctgttaa gggttaaagct 720
gctgaggatt tatgggggac ggattcagat ttatggtttg gatatacaca gcaatcgcac 780
tggcagatgt ttaatggaaa aaactctcgt ccttttagag tacatgacta ccagccagag 840
attttcttaa ctcaacctgt atactcagac ttaccatggg atggcaaagt ccgcatgatt 900
ggcatgggtg cggtagatca ttccaatggg gaaagtgcc aactgtctcg ctcatggaat 960
cgtgcttatt tgatggcagg catggaatgg aaaaacctga ctgtcatgcc acgcatttgg 1020
gggcgtatct ttaaagaggg tagtggcagc cagccagatg ataatcctga tatcttggac 1080
tattatgggt atggtgatgt gcgtttttta tatcaactag aaaataaaaag taatatttca 1140
ggtacgggtac gctataatcc acgctcaggc aaagggtcgt tgcaacttga ctatgtctat 1200
ccgcttggtg aggggaattag tggctatgtt caaatatttc aaggctatgg gcagtctttg 1260
attgattata atcatgaggc gacaagcttt ggcgtcggac ttatgcttaa cgactggatg 1320
ggtctataa 1329

```

<210> 8

<211> 442

<212> PRT

<213> *Moraxella catarrhalis*

<400> 8

```

Met Lys Val Ser Leu Ser Thr Leu Thr Leu Ser Ile Leu Pro Cys Phe
  1          5          10          15
Ala Ile Leu Ala Ile Gln Gln Ala Gln Ala Val Pro Asn Pro Val Ala
  20          25          30
Phe Val Asp Glu Val Arg Ser Glu Asn Asp Leu Gly Gln Asp Asn Glu
  35          40          45
Leu Pro Ile Asp Val Gln Ser Ala Thr Gln Ser Ala Ser Thr Asp Thr
  50          55          60
Ala Asn Pro Leu Asp Glu His Glu Pro Glu Leu Tyr Thr Thr Ala Leu
  65          70          75          80
Glu Asn Lys Thr Met Leu Ile Asn Cys Ser Ala Leu Asn Gln Asp Ile
  85          90          95
Met Arg Leu Ala Cys Tyr Asp Thr Leu Val His Gly Glu Thr Pro Ala
 100          105          110
Val Ile Lys Thr Lys Arg Ser Ile Arg Leu Asp Glu Thr Ile Trp Gln
 115          120          125
Thr Ile Lys Gly Lys Pro Gln Val Val Tyr Gln Glu Thr Thr Asp Pro
 130          135          140
Ile Phe Leu Met Gly Asn Glu Lys Gly Met Leu Thr Lys Lys Asp Ala
 145          150          155          160
Lys Gln Leu Glu Tyr Ala Ala Lys Gln Phe Thr Pro Leu Ser Leu Ser
 165          170          175
Phe Asp Leu Asp Arg Asn Asn Thr Pro Leu Trp Ser Ser Arg Pro His
 180          185          190
Asn Pro Met Tyr Val Leu Pro Ile Phe Met His Gly Lys Pro Asn Arg
 195          200          205
Ser Pro Asn Thr Pro Ser His Glu Ala Lys Gln Phe Thr Pro Asn Glu
 210          215          220
Phe Arg Ala Pro Glu Leu Lys Phe Gln Val Ser Val Lys Val Lys Ala
 225          230          235          240
Ala Glu Asp Leu Trp Gly Thr Asp Ser Asp Leu Trp Phe Gly Tyr Thr
 245          250          255
Gln Gln Ser His Trp Gln Ile Phe Asn Gly Lys Asn Ser Arg Pro Phe
 260          265          270
Arg Val His Asp Tyr Gln Pro Glu Ile Phe Leu Thr Gln Pro Val Tyr
 275          280          285
Ser Asp Leu Pro Trp Asp Gly Lys Val Arg Met Ile Gly Met Gly Ala

```

```

      290              295              300
Val His His Ser Asn Gly Glu Ser Ala Lys Leu Ser Arg Ser Trp Asn
305              310              315              320
Arg Ala Tyr Leu Met Ala Gly Met Glu Trp Lys Asn Leu Thr Val Met
              325              330              335
Pro Arg Ile Trp Gly Arg Ile Phe Lys Glu Gly Ser Gly Ser Gln Pro
              340              345              350
Asp Asp Asn Pro Asp Ile Leu Asp Tyr Tyr Gly Tyr Gly Asp Val Arg
              355              360              365
Phe Leu Tyr Gln Leu Glu Asn Lys Ser Asn Ile Ser Gly Thr Val Arg
              370              375              380
Tyr Asn Pro Arg Ser Gly Lys Gly Ala Leu Gln Leu Asp Tyr Val Tyr
385              390              395              400
Pro Leu Gly Lys Gly Ile Ser Gly Tyr Phe Gln Ile Phe Gln Gly Tyr
              405              410              415
Gly Gln Ser Leu Ile Asp Tyr Asn His Glu Ala Thr Ser Phe Gly Val
              420              425              430
Gly Leu Met Leu Asn Asp Trp Met Gly Leu
              435              440

```

```

<210> 9
<211> 22
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Primer sequence

```

```

<400> 9
gattttaagag tatgttatga tg

```

22

```

<210> 10
<211> 21
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Primer sequence

```

```

<400> 10
gtatggggttg atcaaataca g

```

21

```

<210> 11
<211> 58
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Oligonucleotide

```

```

<400> 11
aagggcccaa ttacgcagag gggatcccaa gctgtaccaa atcctgtggc atttggtg

```

58

```

<210> 12
<211> 60
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Oligonucleotide

```

```

<400> 12

```

aagggcccaa ttacgcagag ggtcgactta ttatagaccc atccagtcgt taagcataag 60

<210> 13

<211> 1000

<212> DNA

<213> Moraxella catarrhalis

<400> 13

acttggcgaa	aataccattt	atatcgattg	tgatgttata	caggcagatg	gcggtacacg	60
cacagccagt	atcagtgggtg	ctgcgggtggc	acttattgat	gctttagaac	acttgcagcg	120
tcgtaaaaag	cttaccceaag	atccgctttt	gggcttggtg	gcagcgggtt	ctgtgggtgt	180
taatcaaggc	cgtgtattgc	ttgatttggg	ttatgctgaa	gattcaactt	gtgataccga	240
tttaaattgtg	gtcatgacgc	aggcagggtg	gtttattgag	attcaaggca	cagcagaaga	300
aaagccattt	actcgtgctg	aagctaattgc	gatgcttgat	ttggcagagc	tgggaattgg	360
gcagattatc	gaagcccaaa	agcaagtatt	aggctggtga	tatgctaata	ggtgaagata	420
atggcgtgat	catcacatta	aatggacaag	taaaagaccc	attatttttg	tggtcgatga	480
tattgctgct	gctgggtgtc	ttgggtggcaa	tcatttggtt	gattgcaccc	gttttttatg	540
caatcgggtgc	gttggcttta	tttgcagttg	tggtatttgt	gtttaatat	caaaggcaaa	600
aagccaaaac	ttgtcatatg	ttttcacaag	gtcgttgtaa	gattacgtcc	aaacgctttg	660
agattcataa	caaataccta	accttatcag	catcggcaac	aatatctgct	aaagataaca	720
aaatgacaat	tggtgatcgg	ggcattgaat	atcattttac	agggttttgct	gatgaccgtg	780
aaattaatat	agccaaacag	gtacttttgg	gaaagtcaat	caaaaccaat	gcggtggcgg	840
taacattggc	taagtagttg	ttgtgataca	gacaggttgg	atggtcttta	actccacca	900
cctaactttt	tctttgtttg	gatttaagag	tatgttatga	tgggcaggat	tttattttta	960
gtcatcatth	aatgcaatca	gttgctcaga	gtagccgttc			1000